

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Level and GCE Advanced Subsidiary Level

MARK SCHEME for the May/June 2006 question paper

9701 CHEMISTRY

9701/02

Paper 2

Maximum raw mark 60

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

The minimum marks in these components needed for various grades were previously published with these mark schemes, but are now instead included in the Report on the Examination for this session.

- CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2006 question papers for most IGCSE and GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 1	Mark Scheme	Syllabus	Paper
	GCE A/AS Level – May/June 2006	9701	02

- 1 (a) (i) ammonia/ NH_3 (1)
- (ii) NH_4^+ (1)
- (iii) iron(II) hydroxide/ $\text{Fe}(\text{OH})_2$ (1) [3]
- (b) barium sulphate/ BaSO_4 (1) [1]
- (c) (i) FeSO_4 (1)
- $(\text{NH}_4)_2\text{SO}_4$ (1)
- (ii) $\text{FeSO}_4 = 151.9$ (allow 151.8 to 152) (1)
- $(\text{NH}_4)_2\text{SO}_4 = 132.1$ (allow 132) (1)
- (iii) $x\text{H}_2\text{O} = 392 - (132.1 + 151.9) = 108$ (1)
- $x = \frac{108}{18} = 6$ (1)
- allow e.c.f. on candidate's sulphate in (c)(i)
e.c.f. answer must be a whole number [6]

[Total: 10]

Page 2	Mark Scheme	Syllabus	Paper
	GCE A/AS Level – May/June 2006	9701	02

- 2 (a)
$$\begin{array}{c} \times \\ \circ \\ \text{H}^{\times} \text{C}^{\times} \text{C}^{\circ} \text{H} \\ \times \\ \circ \end{array}$$
 (1) [1]
- (b) $n = \frac{PV}{RT} = \frac{(1515 \times 10^3) \times (76 \times 10^{-3})}{8.31 \times 298}$ (1)
- = 46.5 (1) [2]
- (c) (i) $\text{CaC}_2 + 2\text{H}_2\text{O} \rightarrow \text{Ca(OH)}_2 + \text{C}_2\text{H}_2$ (1)
- (ii) $n(\text{C}_2\text{H}_2) = n(\text{CaC}_2) = 100 \times 46.5$ (1)
- mass of $\text{CaC}_2 = 100 \times 46.5 \times 64 =$
- = 297 570 g
- = 297.6 kg (accept 298 kg)
- correct units necessary (1)
- allow e.c.f. on candidate's answer in (b) [3]
- (d) $\text{C}_2\text{H}_2(\text{g}) + \frac{5}{2}\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{g})$
- bonds broken: $2(\text{H-C}) \quad 2 \times 410 \quad = \quad 820$
- $\text{C}\equiv\text{C} \quad 840 \quad = \quad 840$
- $\frac{5}{2}(\text{O}=\text{O}) \quad \frac{5}{2} \times 496 \quad = \quad \underline{1240}$
- 2900 kJ mol⁻¹ (1)
- bonds made: $4(\text{C}=\text{O}) \quad 4 \times 740 \quad = \quad 2960$
- $2(\text{O}-\text{H}) \quad 2 \times 460 \quad = \quad \underline{920}$
- 3880 kJ mol⁻¹ (1)
- $\Delta H_{\text{comb}} = -3880 + 2900 = -980 \text{ kJ mol}^{-1}$ (1)
- allow e.c.f. on incorrect bonds made/broken [3]
- (e) (i) the enthalpy/energy change when one mole of a substance (1)
- is burned in an excess of air/oxygen
- or completely combusted
- under standard conditions (1)
- (ii) calculation in (d) includes $\text{H}_2\text{O}(\text{g})$ whereas ΔH_{comb} involves $\text{H}_2\text{O}(\text{l})$
- or average bond energy terms are used in the *Data Booklet* (1) [3]

[Total: 12]

Page 3	Mark Scheme	Syllabus	Paper
	GCE A/AS Level – May/June 2006	9701	02

3 (a)

halogen	colour	physical state at room temperature
chlorine	green/yellow	gas
bromine	orange/red	liquid
iodine	black	solid

three colours correct (1)

three states correct (1) [2]

(b) (i) $MgCl_2$ white fumes/steamy fumes (of HCl) (1)

$MgBr_2$ red colour (of Br_2) or steamy fumes (of HBr) (1)

MgI_2 purple colour (of I_2) or black solid (I_2) or yellow solid (S) or stinking gas (H_2S) (1)

(ii) $MgCl_2 + H_2SO_4 \rightarrow MgSO_4 + 2HCl$
allow
 $MgCl_2 + 2H_2SO_4 \rightarrow Mg(HSO_4)_2 + 2HCl$ (1) [4]

(c) (i) $AgBr$ /silver bromide (1)

(ii) $AgBr(s) + 2NH_3(aq) \rightarrow Ag(NH_3)_2Br(aq)$
equation (may be ionic) (1)

state symbols (1)

allow ecf on wrong halide in (i) [3]

(d) (i) HCl no reaction (1)

HI purple vapour/black solid (1)

$2HI \rightarrow H_2 + I_2$ (1)

(ii) bond energy in HCl is high (1)

bond energy in HI is lower/more easily broken (1)

(iii) hot glass rod provides activation energy (1) [6]

[Total: 15]

Page 4	Mark Scheme	Syllabus	Paper
	GCE A/AS Level – May/June 2006	9701	02

- 4 (a) (i) $\text{BrCH}_2\text{CHBrCH}_2\text{OH}$ (1)
- (ii) $\text{CH}_2=\text{CHCO}_2\text{H}$ (1) [2]
- (b) oxidation (1) [1]
- (c) structural **or** functional group isomerism (1) [1]
- (d) **step I**
 $\text{CH}_2=\text{CHCH}_2\text{OH} \rightarrow \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
- H_2 (1)
- Ni catalyst and heat **or** Pt at room temperature (1)
- step II** $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} \rightarrow \text{CH}_3\text{CH}_2\text{CHO}$
- acidified dichromate(VI) (1)
- heat **not** under reflux
or distil off aldehyde as it is formed (1)
- for both steps, conditions mark is only awarded if correct reagent is given [4]
- (e) **both** oxidation and reduction have occurred (1) [1]
- (f) (i) $\text{HOCH}_2\text{CHOHCH}_2\text{OH}$ formed
- $\text{CH}_2=\text{CH}-$ forms $\text{HOCH}_2\text{CHOH}-$ (1)
- $-\text{CH}_2\text{OH}$ is unchanged (1)
- (ii) $\text{HO}_2\text{CCO}_2\text{H}$ (1) [3]
- [Total: 12]

Page 5	Mark Scheme	Syllabus	Paper
	GCE A/AS Level – May/June 2006	9701	02

- 5 (a) A reaction in which one atom or group of atoms replaces another. (1) [1]
- (b) (i) $C_2H_5Br + NaOH \rightarrow C_2H_5OH + NaBr$
allow OH^- (1)
- (ii) heat with aqueous $NaOH/OH^-$ (1) [2]
- (c) (i) $C_2H_5Br + KCN \rightarrow C_2H_5CN + KBr$
allow CN^- or $NaCN$ (1)
- (ii) heat with KCN under reflux in ethanol (1) [2]
- (d) (i) K is $C_2H_5Cl/C_2H_5Br/C_2H_5I$ (1)
L is C_2H_5CN (1)
- (ii) **step I**
reagent - PCl_3 or PCl_5 or $SOCl_2$
or $NaBr + conc\ H_2SO_4$ or $P + Br_2$
or $P + I_2$ (1)
conditions - room temperature or heat under reflux depending on reagent (1)
- step III**
reagent - mineral acid (1)
conditions - heat under reflux (1)
- for both steps, conditions mark is only awarded if correct reagent is given [6]
- [Total: 11]