

# Mark Scheme (Results)

Summer 2016

Pearson Edexcel GCE  
in Chemistry (6CH01) Paper 01  
The Core Principles of Chemistry

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2016

Publications Code 46656\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2016

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
  - i) ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear
  - ii) select and use a form and style of writing appropriate to purpose and to complex subject matter
  - iii) organise information clearly and coherently, using specialist vocabulary when appropriate

## Using the Mark Scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.

( ) means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer.

ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

### Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities.

Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

**Section A (multiple choice)**

Question Number	Correct Answer	Reject	Mark
<b>1</b>	B		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>2</b>	C		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>3</b>	D		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>4</b>	D		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>5</b>	B		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>6</b>	B		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>7</b>	B		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>8a</b>	A		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>8b</b>	B		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>8c</b>	D		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>9</b>	D		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>10</b>	A		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>11</b>	A		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>12</b>	A		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>13</b>	D		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>14</b>	C		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>15</b>	D		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>16a</b>	C		<b>(1)</b>



Question Number	Correct Answer	Reject	Mark
<b>16b</b>	C		<b>(1)</b>

Question Number	Correct Answer	Reject	Mark
<b>17</b>	B		<b>(1)</b>

## Section B

Question Number	Acceptable Answers	Reject	Mark
<b>18a(i)</b>	<p>Any two of  <math>O^+</math>, <math>O^{2+}</math>, <math>O_2^+</math>, <math>O_2^{2+}</math>            (1) for each correct ion</p> <p>ALLOW  <math>^{16}O^+</math>, <math>^{16}O^{2+}</math>, <math>(^{16}O)_2^+</math>, <math>(^{16}O)_2^{2+}</math>  <math>^{16}O_2^+</math>, <math>^{16}O_2^{2+}</math></p> <p><math>O=O^+</math> / <math>O=O^{2+}</math> for <math>O_2</math> ions</p> <p>Added mass numbers which describe a diatomic ion eg <math>^{32}O_2^+</math></p> <p>Added round or square brackets</p>	<p><math>O^-</math>  <math>O^{2-}</math>            Ions of <math>O_3</math></p> <p>Incorrect mass numbers eg  <math>^{32}O^+</math></p> <p>Added incorrect atomic numbers            Eg <math>^{16}_9O^+</math></p>	<b>(2)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>18a(ii)</b>	<p>The magnetic field/            electromagnet/ electromagnetic field            OR            Deflection by magnetic field</p> <p>ALLOW            Deflection and magnetic field</p>	<p>Gravitational field</p> <p>Just deflector/deflection</p> <p>Electric field</p> <p>Vacuum and magnetic field</p> <p>Detector/ detection</p>	<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
18a(iii)	<p><b>Two curved</b> lines going towards the detector region with at least one hitting the detector</p> <p>ALLOW Section of straight line before curve starts if magnetic field position is not shown Line may go up very slightly before it curves down, probably to keep it clear of lower line.</p>  <p style="text-align: right;">(1)</p> <p>Labelling of paths depends on ions chosen:</p> <p><b>Heavier ion</b> shown as <b>less deflected</b> OR <math>O^{2+}</math> more deflected than <math>O_2^+</math> OR Ion with <b>lower charge</b> shown as <b>less deflected</b></p> <p>ALLOW Ions with negative charges (as already penalised in (i)) (1)</p> <p>If chosen ions are <math>O^+</math> and <math>O_2^{2+}</math> they will not be separated – answer must make this clear</p> 	<p>Straight lines Curvature away from detector/ concave curvature</p> <p>Line turning back upwards</p> <p>Species which are not ions of oxygen</p>	<b>(2)</b>



Question Number	Acceptable Answers	Reject	Mark
<b>18(b)</b>	<p><b>Look at final answer</b>  <b>16.004 scores (2)</b>  <b>16.00445 scores (1)</b></p> <p>Correct expression with incorrect final answer scores (1)</p> <p><math>(16 \times 99.759 + 17 \times 0.037 + 18 \times 0.204) / 100</math>  OR  <math>(16 \times 0.99759 + 17 \times 0.00037 + 18 \times 0.00204)</math> (1)</p> <p>= 16.00445  = 16.004 (1)</p> <p>Ignore units</p>	16.005	<b>(2)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>18(c)</b>	<p>Isotopic composition of oxygen in air varies</p> <p>ALLOW  The abundance of the isotopes of oxygen varies</p> <p>OR  Oxygen standard was introduced before existence of oxygen isotopes was known</p> <p>OR  Some scientists used a standard based on one isotope while others used a value based on mixture in natural abundance</p> <p>OR  The answer is inaccurate unless a specified isotope is used</p> <p>OR  <math>^{12}\text{C}</math> standard used because there are many <math>^{12}\text{C}</math> compounds which can be used to calibrate the mass spectrometer</p> <p>ALLOW  It was difficult to obtain pure oxygen from air.</p>	<p>Air contains other gases</p> <p>Air contains many isotopes</p> <p>Oxygen has many isotopes</p> <p>Just <math>^{12}\text{C}</math> standard is better'  <math>^{12}\text{C}</math> standard gives a whole number</p>	<b>(1)</b>


Question Number	Acceptable Answers	Reject	Mark
<b>18(d)</b>	No difference as both isotopes have the same number of protons (and electrons)/ the same nuclear charge  IGNORE Same electronic configuration  OR No difference as <b>only</b> number of neutrons is different		<b>(1)</b>

**(Total for Question 18 = 9 marks)**

Question Number	Acceptable Answers	Reject	Mark
<b>19(a)</b>	$\text{Mg(g)} \rightarrow \text{Mg}^+(\text{g}) + \text{e}^{(-)}$ ALLOW $\text{Mg(g)} - \text{e}^{(-)} \rightarrow \text{Mg}^+(\text{g})$ Loss of electron to form $\text{Mg}^+$ (1) IGNORE (g) sign on electron State symbols ALLOW Provided the equation involves magnesium, even if electron is added to the wrong side. (1)	Formation of $\text{Mg}^{2+}$	<b>(2)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19(b)</b>	$(1s^2) 2s^2 2p^6 3s^2 3p^1$ ALLOW Capital s and/or p, subscripts $2p_x^2 2p_y^2 2p_z^2 3p_x^1$ $3p_y^1 / 3p_z^1$ for $3p_x^1$		<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
*19(c)(i)	<p><b>MP1</b> Mg to Al: Electron removed from Al is from a higher energy level (3p rather than 3s) ALLOW Electron removed in Al is (more) shielded (by 3s) IGNORE Outer electron is further from nucleus Full sub-shell is more stable than part filled sub-shell (1)</p> <p><b>MP2</b> Al to Si: Si has one more proton than Al/ has greater nuclear charge, <b>and</b> electrons removed in both cases are 3p / same sub-shell / are equally shielded (1)</p> <p><b>MP3</b> EITHER The attraction of the extra proton in Al is less than the effect of the higher energy level/ the shielding</p> <p>OR Electron removed from Si is closer to nucleus (than Al) ALLOW Silicon is smaller in size (1)</p>		<b>(3)</b>

Question Number	Acceptable Answers	Reject	Mark
19(c)(ii)	 <p data-bbox="400 1010 1163 1823"> <b>MP1</b>            S does not follow trend            (P is above Si followed by dip in graph from P to S rising again to Cl and Ar) (1)         </p> <p data-bbox="400 1189 1163 1435"> <b>MP2</b>            S has one (3)p orbital which has two electrons/ paired electrons/ is fully occupied            OR            S has <math>3p_x^2, 3p_y^1, 3p_z^1</math>            OR            Electron in box diagram for S         </p> <p data-bbox="400 1469 1163 1541">           ALLOW            S has <b>a pair of</b> electrons in the (3)p subshell (1)         </p> <p data-bbox="400 1581 1163 1823"> <b>MP3</b>            A paired electron is easier to remove            OR            paired electrons repel each other            ALLOW            half filled sub-shell (in P) is stable (1)         </p>	<p data-bbox="1190 1227 1299 1335">Just "S has <math>3p^4</math>"</p> <p data-bbox="1190 1368 1299 1440">d orbital</p> <p data-bbox="1190 1688 1299 1823">P has <b>a</b> half filled orbital</p>	(3)

Question Number	Acceptable Answers	Reject	Mark
<b>19(d)</b>	<p>Four x round Si sharing one • with each Cl (1)            Seven • round each Cl sharing one x with each Si (1)</p> <p style="text-align: center;"> <math display="block">\begin{array}{c} \cdot\cdot \\ : \text{Cl} : \\ \cdot \times \end{array}</math> <math display="block">\begin{array}{c} \cdot\cdot \quad \times \quad \cdot\cdot \\ : \text{Cl} \cdot \text{Si} \cdot \text{Cl} : \\ \cdot\cdot \quad \cdot\cdot \end{array}</math> <math display="block">\begin{array}{c} \times \cdot \\ : \text{Cl} : \\ \cdot\cdot \end{array}</math> </p> <p>ALLOW Reversed symbols</p>		<b>(2)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>*19(e) (i)</b>	<p><b>MP1</b>  <math>\text{I}^-</math> / anion becomes distorted / not spherical. May be shown in a diagram (1)</p> <p><b>MP2</b>  <math>\text{Mg}^{2+}</math> has high(er) charge <b>and</b> small(er) radius/ <math>\text{Mg}^{2+}</math> has high charge density (1)</p> <p><b>MP3</b>            Bonding in magnesium iodide has some covalent character</p> <p>OR            Orbitals of <math>\text{Mg}^{2+}</math> and <math>\text{I}^-</math> overlap/ <math>\text{Mg}^{2+}</math> shares some of the <math>\text{I}^-</math> electrons</p> <p>OR  <math>\text{Mg}^{2+}</math> and <math>\text{I}^-</math> ions are not completely separate (1)</p>	<p>Iodine becomes distorted            Just "electrons in outer shell are attracted"</p> <p>Atoms of Mg have a small (atomic) radius</p>	<b>(3)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>19(e)(ii)</b>	<p>Experimental/ Born Haber cycle and theoretical/ calculated lattice energies are <b>different</b></p> <p>OR</p> <p>Experimental/ Born Haber cycle lattice energy is more exothermic/ more negative than theoretical/ calculated lattice energy</p> <p>ALLOW</p> <p>Greater for more negative</p> <p>IGNORE</p> <p>Comments about melting temperature</p>	<p>Just "Compare Experimental/ Born Haber cycle and theoretical/ calculated lattice energies"</p> <p>Use of electron density map</p>	<b>(1)</b>

**(Total for Question 19 = 15 marks)**

Question Number	Acceptable Answers	Reject	Mark
<b>20(a) (i)</b>	(Different) boiling temperatures/ boiling points  ALLOW Range of boiling temperatures		<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20(a) (ii)</b>	<b>Cracking:</b> breaking of carbon chain (in a hydrocarbon/ alkane) to give shorter chain hydrocarbon(s)/ smaller molecules  OR breaking a hydrocarbon/ alkane to give smaller molecules  OR Breaking an alkane to give an alkene <b>and</b> (a smaller) alkane/ hydrogen (1)  <b>Reforming:</b> converting straight chain to a (more) branched chain/ ring/ arene / aromatic compound  ALLOW Specific examples (1)  IGNORE Makes more useful compounds Converting low octane (fuels) into high octane (fuels)	Just "Breaking a hydrocarbon"  Just "Breaking a molecule"  Breaking a hydrocarbon to form branched chains or ring structures	<b>(2)</b>



Question Number	Acceptable Answers	Reject	Mark
<b>20(a) (iii)</b>	<p>Look at final answer:  <b>+71</b> (kJ mol<sup>-1</sup>) scores <b>3</b> marks  <b>-71/ 71</b> (kJ mol<sup>-1</sup>) scores <b>2</b> marks  <b>-5825</b> (kJ mol<sup>-1</sup>) scores <b>1</b> mark</p> <p>Method:</p> $  \begin{array}{c}  \text{C}_4\text{H}_{10} \rightarrow \text{C}_3\text{H}_6 + \text{CH}_4 \\  (+13/2 \text{ O}_2) \qquad \qquad \qquad (+13/2 \text{ O}_2) \\  \swarrow \qquad \qquad \searrow \\  -2877 \qquad \qquad -2058-890 \text{ /-2948} \\  \qquad \qquad \qquad \downarrow \\  \qquad \qquad \qquad 4\text{CO}_2 + 5\text{H}_2\text{O}  \end{array}  $ <p><b>MP1</b>  Labelled cycle  OR  use of  <math>\Delta H = \sum \Delta H_{\text{combustion}} \text{ reactants} - \sum \Delta H_{\text{combustion}} \text{ products}</math> (1)</p> <p><b>MP2</b>  <math>\Delta H = (-2877 - (-2058 + (-890)))</math> (1)</p> <p><b>MP3</b>  = +71 (kJ mol<sup>-1</sup>) (1)</p>	Incorrect units	<b>(3)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20(a) (iv)</b>	$\text{C}_4\text{H}_{10} \rightarrow \text{C}_2\text{H}_6 + \text{C}_2\text{H}_4$ OR $\text{C}_4\text{H}_{10} \rightarrow \text{C}_4\text{H}_8 + \text{H}_2$ OR $\text{C}_4\text{H}_{10} \rightarrow 2\text{C}_2\text{H}_4 + \text{H}_2$ <p>ALLOW  Breakdown of multiple butanes</p> <p>Ignore state symbols, even if incorrect</p>	$\text{C}_4\text{H}_{10} \rightarrow \text{C}_3\text{H}_6 + \text{CH}_4$  Charged products eg $\text{C}_2\text{H}_5^+$  Free radicals eg $\text{C}_2\text{H}_5^\cdot$	<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20b(i)</b>	<p>Look at final answer:  <b>-2050</b> (kJ mol<sup>-1</sup>) or anything correctly rounded from -2046.528  <b>(-2047, -2046.5, -2046.53)</b>  scores <b>3</b> marks</p> <p><b>+2050/ 2050</b> (kJ mol<sup>-1</sup>) scores <b>2</b> marks</p> <p>Incorrect rounding scores 2 marks</p> <p>Correct value without sign scores 2 marks</p> <p>Energy transferred = (200 x 4.18 x 34.0)  =28424 (J)  IGNORE  Sign if given (1)</p> <p>Mol pentane =(1.0/72) = 0.01389 /  0.0139 (1)</p> <p><math>\Delta H = - (-28424 \div (1/72 \times 1000))</math>  = -2046.528 (kJ mol<sup>-1</sup>)</p> <p>ALLOW  TE from MP 1 and 2 provided moles of pentane is not taken as 1 (1)</p> <p>NOTE  Use of 0.0139 mol gives  -2044.9 (kJ mol<sup>-1</sup>) giving 3 marks  Use of 0.0138 mol gives  -2059.7 (kJ mol<sup>-1</sup>) giving 2 marks  Use of 0.014 mol gives  -2030.29 (kJ mol<sup>-1</sup>) giving 2 marks</p> <p>Ignore SF except one or two</p>		<b>(3)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20(b) (ii)</b>	Incomplete combustion OR Loss of pentane by evaporation  ALLOW Volume of water too large to heat evenly Water not stirred evenly Small change in mass inaccurate Heat capacity of /energy needed to heat calorimeter not included	Incomplete reaction Loss of water by evaporation  Heat losses  Conditions not standard  Measuring errors  Pentane impure	<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20(b) (iii)</b>	Pentane is <b>very</b> volatile/ has low boiling temperature so risk of explosion  OR Has <b>high</b> flammability  IGNORE Reaction is very exothermic	Just "it is flammable"  Vapour is toxic Combustion products/ CO toxic	<b>(1)</b>


Question Number	Acceptable Answers	Reject	Mark
<b>20(c) (i)</b>	$C_5H_{12} + 8O_2 \rightarrow 5CO_2 + 6H_2O$ Allow multiples  Ignore state symbols even if incorrect		<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20(c)(ii)</b>	<p>Bonds broken are four C-C twelve C-H eight O=O (1)</p> <p>Bonds made are ten C=O twelve O-H (1)</p> <p>ALLOW TE from (c)(i)</p> <p>If all five bonds are named but formulae not given eg oxygen-oxygen bonds, max 1</p> <p>If all five bonds are correctly identified by formula but numbers are incorrect or missing, max 1</p>	<p>O-O single bonds</p> <p>C-O single bonds</p>	<b>(2)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>20(c)(iii)</b>	<p>The (total) bond energy of the bonds formed is greater than the bond energy of the bonds broken</p> <p>OR</p> <p>Energy released forming new bonds &gt; energy needed to break old bonds</p> <p>OR</p> <p>The sum of the bond energies of the products is greater than the sum of the bond energies of the reactants.</p>	<p>Just "more bonds are made than broken"</p> <p>Answers referring to energy needed to make bonds</p> <p>Energy contained by bonds in reactants &gt; energy contained by bonds in products</p>	<b>(1)</b>

**(Total for question 20 = 16 marks)**

Question Number	Acceptable Answers	Reject	Mark
<b>21(a)(i)</b>	Species/ atom/ molecule/ particle with an <b>unpaired electron</b>  ALLOW An element with an <b>unpaired electron</b>  IGNORE Reference to neutral species /lack of charge	Just "with a single electron"  A lone electron  Charged particle with an unpaired electron	<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>21(a)(ii)</b>	  Half arrows going from bond to Cl or just beyond <b>and</b> product $2\text{Cl}\cdot$ / $\text{Cl}\cdot + \text{Cl}\cdot$	Cl without •	<b>(1)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>21a(iii)</b>	$\text{C}_2\text{H}_6 + \text{Cl}\cdot \rightarrow \text{C}_2\text{H}_5\cdot + \text{HCl}$ ALLOW Structural formulae e.g. $\text{CH}_3\text{CH}_3$ OR displayed  IGNORE Production of $\text{C}_2\text{H}_5\text{Cl}$ from $\text{C}_2\text{H}_5\cdot$ if first step is correct (1)  Propagation (1)  The second mark is independent of the first	$\text{C}_2\text{H}_5^+$	<b>(2)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>21a(iv)</b>	$C_2H_5\cdot + C_2H_5\cdot \rightarrow C_4H_{10}$  ALLOW Structural formulae e.g. $CH_3CH_2\cdot$ / $\cdot CH_3CH_2$ OR displayed  IGNORE $Cl\cdot + Cl\cdot \rightarrow Cl_2$	Methyl or propyl radicals	<b>(1)</b>


Question Number	Acceptable Answers	Reject	Mark
<b>21b(i)</b>	<p> <math>\sigma</math> bond between C atoms shown as 2 overlapping orbitals/ one electron cloud/ single bond (1)         </p> <p> <math>\pi</math> bond above and below <math>\sigma</math> bond shown as two electron clouds/ overlapping p orbitals/ p orbitals linked by a line / a curved line above and below single bond (1)         </p> <p>Both bonds must be labelled for 2 marks.</p>		<b>(2)</b>


Question Number	Acceptable Answers	Reject	Mark
<b>*21b(ii)</b>	<p><b>MP1</b>  <math>\sigma</math> bond remains  ALLOW  The product contains <math>\sigma</math> bonds only  (1)</p> <p><b>MP2</b>  <math>\pi</math> bonds break because they are weaker (than <math>\sigma</math> bonds)  ALLOW  <math>\pi</math> bonds break because <math>\sigma</math> bonds are stronger  (1)</p> <p><b>MP3</b>  Breaking the <math>\pi</math> bond results in carbocation intermediate / positively charged carbon forming</p> <p>OR  <math>\pi</math> orbital overlap is lateral/ sideways /between parallel orbitals (making <math>\pi</math> bonds break/ weak)</p> <p>OR  The <math>\sigma</math> bonds are much stronger (than the <math>\pi</math> bond) because of more effective (orbital) overlap  (1)</p>		<b>(3)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>21(b)(iii)</b>	<p>From: Purple/ pink (solution)  To: colourless  (1)</p> $\begin{array}{ccccccc} & & \text{H} & & \text{H} & & \\ & &   & &   & & \\ \text{H} & - & \text{O} & - & \text{C} & - & \text{C} & - & \text{O} & - & \text{H} \\ & &   & &   & & \\ & & \text{H} & & \text{H} & & \end{array}$ <p>(1)</p> <p>Any orientation  Don't penalise undisplayed OH</p> <p>Don't penalise bonds going to middle of undisplayed OH</p>	<p>To brown</p> <p>Molecular/  structural/  skeletal formulae</p> <p>C bonded to H  of OH</p>	<b>(2)</b>

Question Number	Acceptable Answers	Reject	Mark
<b>21 (b) (iv)</b>	<p>Second mark depends on use of bromine/ solution of bromine for test.</p> <p>EITHER            Test: add bromine water / Br<sub>2</sub>(aq)            ALLOW            Add bromine in organic solvent/            bromine dissolved in hexane/            bromine in 1,1,1-trichloroethane            (1)</p> <p>From: brown/ red-brown/orange/            yellow            To: colourless (1)</p> <p>OR            Add bromine / Br<sub>2</sub>            (1)</p> <p>From: brown/ red-brown            To: colourless (1)</p>		<b>(2)</b>



Question Number	Acceptable Answers	Reject	Mark
21(b)(v)	 <p>Dipole on HBr (1)</p> <p>Curly arrow from C=C double bond to H<sup>δ+</sup> of HBr <b>and</b> curly arrow from H-Br bond to Br (1)</p> <p>Correct intermediate with + charge (1)</p> <p>Curly arrow from Br<sup>-</sup> to C<sup>+</sup> and formula of product</p> <p>ALLOW Curly arrow from anywhere on Br, including the - sign or lone pair (which is optional) (1)</p>	Half arrows	(4)

Question Number	Acceptable Answers	Reject	Mark
21(c)	 <p>(1)</p> <p>Suitable catalyst nickel/ platinum/ palladium (1)</p> <p>Ignore references to temperature, pressure, uv light</p>	Use of H, H <sup>+</sup>  Zeolite catalyst	(2)

(Total for Question 21 = 20 marks)

TOTAL FOR PAPER = 80 MARKS

