

Mark Scheme (Results)

October 2020

Pearson Edexcel International Advanced Subsidiary Level In Chemistry (WCH13)

Paper 1: Practical Skills in Chemistry I

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2020
Publications Code WCH13_01_2010_MS
All the material in this publication is copyright
© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Using the mark scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit. () means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer. ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities. Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Question	Answer		Additional Guidance	Mark
Number				
1(a)	An answer that makes reference to the following points:			(2)
	 nitrogen dioxide / NO₂ / nitrogen(IV) oxide oxygen / O₂ 	(1) (1)	Allow N ₂ O ₄ / dinitrogen tetroxide Do not award O	
			If name and formula are given both must be correct Ignore reference to observations, even if incorrect	

Question Number	Answer	Additional Guidance	Mark
1(b)	 nitrate / NO₃⁻ / nitrate(V) 	Do not award nitrate(III) or nitrite	(1)
		If name and formula are given both must be correct	

Question Number		Additional Guidance	Mark
1(c)	• barium (ion) /Ba ²⁺	Do not award Ba / Ba ⁺	(1)
		If name and formula given both must be correct	

Question	Answer		Additional Guidance	Mark
Number				
1(d)	An answer that makes reference to the following points:		TE for both marks on 1(c) for incorrect group 2 / group 1 cation	(2)
	• (Solid A) Ba(NO ₃) ₂	(1)	If only names given penalise once only	
	• BaO	(1)	Ignore state symbols, even if incorrect	

Question	Answer		Additional Guidance	Mark
Number				
1(e)	An answer that makes reference to the following points:		TE for both marks on 1(c) for incorrect group 2	(2)
			/ group 1 cation	
	(Sodium hydroxide)			
	 no change / no reaction / colourless solution 	(1)	Allow (white) precipitate does not form	
			Ignore nothing / no observation	
			Ignore just colourless / remains clear	
			Ignore solid dissolves	
			Do not award white solution	
			Do not award effervescence / gas evolved	
	(Sulfuric acid)		Do not award enervescence / gas evolved	
	white and precipitate	(1)	Allow ppt(e)/solid/crystals for precipitate	
	- Write and precipitate	` ,	, · · · · · · · · · · · · · · · · · · ·	
			Ignore name or formula of precipitate	
			Do not award effervescence / gas evolved	

(Total for Question 1 = 8 marks)

Question Number	Answer	Additional Guidance	Mark
2(a)	An answer that makes reference to the following points:	Penalise incorrect formulae once only	(3)
	■ B is silver nitrate (solution) / AgNO ₃	Any 1 correct = 1 mark	
	C is hydrochloric acid / HCl	Any 2 or 3 correct = 2 marks	
	• D is potassium carbonate (solution) / K₂CO₃	4 correct = 3 marks	
	• E is sodium chloride (solution) / NaCl		

Question Number	Answer		Additional Guidance	Mark
2(b)	An answer that makes reference to the following points: Step 1 Improvement (M1) • use (conc.) hydrochloric acid Explanation (M2 – dependent on M1) • (metal) chlorides are more volatile or Improvement (M1) • dip wire in (conc.) hydrochloric acid and place in flame Explanation (M2 – dependent on dipping wire in acid and placing in flame) • to clean the wire / remove traces of previous sample(s) (that might interfere with the result)	(1) (1) (1)	Ignore reference to use of powder/solid Ignore just chlorides give better result Ignore reference to acid strength Ignore nitric acid is oxidising/hazards of nitric acid Ignore reference to NO ₂ (eg toxic/brown) Do not award Cl ⁻ /chloride ions / HCl more volatile Do not award (metal) chlorides are more soluble Ignore dip wire in acid then solid (improvement)to put more sample on wire (explanation)	(6)
	Step 2 Improvement (M3) • use nichrome (wire/loop) / platinum (wire/loop) Explanation (M4 – dependent on use of a different wire) • copper produces a (blue-green) flame colour (so interferes with the result)	(1) (1)	Allow NiCr / Pt Do not award nickel/Ni or chromium/Cr Allow copper will react with nitric acid Allow copper will melt Ignore copper reacts with flame/burns/oxidises Allow nichrome/platinum would not produce a flame colour / are inert/unreactive/do not react Ignore nichrome/platinum less reactive than copper	

Step 3		Allow half-open / should not be closed	
Improvement (M5)		Allow Bunsen with non-luminous flame	
 use the Bunsen burner with the air-hole (fully) open 	(1)	Ignore colour change for flame colour	
 Explanation (M6 – dependent on M5) so flame colour is easier to see (with non-luminous Bunsen flame) / flame is hotter 		Ignore reference to combustion	
raminous Buriser marile, 7 marile is motter		Ignore colour change for flame colour	
or			
 flame colour is more difficult to see (with 			
luminous Bunsen flame when air-hole closed) / flame is too cool / not hot enough	(1)		

(Total for Question 2 = 9 marks)

Question	Answer	Additional Guidance	Mark
Number			
3(a)(i)			(1)
	 hydrogen chloride/HCl((g))/hydrochloric acid/HCl(aq) 	If name and formula are given both must be correct	

Question	Answer	Additional Guidance	Mark
Number			
3(a)(ii)			(1)
	 carbon dioxide / CO₂ 	If name and formula are given both must be correct	

Question Number	Answer		Additional Guidance	Mark
3(a)(iii)	• F = alcohol	(1)	Allow hydroxyl / hydroxy Ignore any reference to primary/secondary/tertiary Do not award hydroxide / OH ⁻	(2)
	• G = carboxylic acid	(1)	Allow carboxylic/carboxyl/carboxy Ignore just acid Ignore hydroxy and carbonyl Do not award carboxic/carbonic	
			Award one mark for both $F = (-)OH$ and $G = (-)COOH / (-)CO2HAward one mark for correct answers reversed$	

Question Number	Answer	Additional Guidance	Mark
3(a)(iv)	H H H – C – C – O – H I I H H (1)	Ignore bond angles and connectivity of OH group unless horizontal C-HO Allow non-displayed OH	(2)
	$H \longrightarrow C \longrightarrow C \longrightarrow C$	Allow non-displayed OH Award 1 mark for two correct skeletal/structural formulae Ignore molecular formulae	

Question Number	Answer	Additional Guidance	Mark
Number 3(a)(v)	An answer that makes reference to the following points: • Yes and F/alcohol would not have a (peak due to) C=O or G/(carboxylic) acid would have a (peak due to) C=O	Allow TE on any alcohol and carboxylic acid in (a)(iv) Allow yes and O-H (peak) values are different in F/alcohol and G/(carboxylic) acid Allow OH / -OH / -O-H for O-H Do not award C-OH for O-H Allow Yes and fingerprint region would be different Ignore just different functional groups give different peaks	(1)
		Ignore just G has an extra peak	

Question	Answer	Additional Guidance	Mark
Number			
3(b)	An answer that makes reference to the following points:	Ignore any reference to miscibility / layers	(4)
	(bromine water)		
	• from brown / orange / yellow (1	Do not award red	
	• to colourless (1	Ignore clear	
	(Benedict's / Fehling's reagent)	Do not award effervescence / gas evolved	
	• from blue (1		
	• to red / brick red and solid / precipitate (1	Allow ppt(e)	
		Allow red-brown for red	
		Do not award brown/orange/pink for red	

(Total for Question 3 = 11 marks)

Question	Answer	Additional Guidance	Mark
Number			
4(a)(i)	 To make sure that all the (hydrochloric) acid / 	Accept completely neutralised/reacted for used	(1)
	HCl((aq)) has been used up	up	
		Ignore 'to ensure the reaction goes to	
		completion'	
		Ignore reference to excess/limiting reagents	
		Ignore reference to temperature change	

Question Number	Answer		Additional Guidance	Mark
4(a)(ii)	An answer that makes reference to the following point	s:		(1)
	 maximum temperature of the mixture: 2 and 	27.5	Allow 27.6	
	temperature change: 6(.	0)	Allow 6.1 (for maximum temperature of 27.6)	

Question Number	Answer	Additional Guidance	Mark
4(b)	An answer that makes reference to the following points:	Example of calculation	(4)
		Ignore SF except 1SF throughout	
	Calculation of energy change (1)	$55 \times 6(.0) \times 4.2 = 1386(J) / 1.386 (kJ)$ TE on temperature change from (a)(ii)	
	Calculation of the moles of HCl (1)	$25 \times 1.0 \div 1000 = 2.5 \times 10^{-2} / 0.025$ (moles) Allow 1/40	
	Calculation of enthalpy change (1)	$1386 \div 0.025 = 55440 \text{ (J mol}^{-1}\text{)}$ TE on energy change TE on moles HCl	
	Negative sign and units (1)	Must follow a calculation where an energy change has been divided by an amount -55440 J mol ⁻⁽¹⁾ / -55.440 kJ mol ⁻⁽¹⁾ Do not award J/mol ⁻⁽¹⁾ or kJ/mol ⁻⁽¹⁾	

Question Number	Answer		Additional Guidance	Mark
4(c)	An answer that makes reference to the following points:			(2)
	less exothermic	(1)	Allow less negative Ignore higher/greater/lower/smaller value	
	glass is a better conductor (of heat)		value	
	or		Allow glass is a worse insulator Allow glass is a good conductor Ignore reference to heat loss	
	polystyrene is a better insulator (of heat)	(1)	Allow polystyrene is a good insulator	
			Allow glass beaker has a higher heat capacity (than polystyrene cup) Allow glass beaker absorbs more heat/energy (than polystyrene cup)	

(Total for Question 4 = 8 marks)

Question	Answer	Additional Guidance	Mark
Number			
5(a)			(1)
	pipette (and filler)	Do not award teat pipette/burette	
	p.pette (and imel)	р.р.столия.	

Question	Answer	Additional Guidance	Mark
Number			
5(b)	• (from) yellow		(2)
	(,	
	• to orange (Allow peach for orange	
		Do not award pink or red	
		Award one mark for correct colours reversed	

Question			Answer			Additional Guidance	Mark
Number							(4)
5(c)(i)	_				_		(1)
		1	2	3		All 3 values correct	
		25.25	25.0(0)	24.8(0)			
	L				1	Do not award 25 for titration number 2	

Question	Answer	Additional Guidance	Mark
Number			
5(c)(ii)	(1) 2 12 1 2 24 0(0) (3)		(1)
	• (titres 2 and 3 chosen) = 24.9(0) (cm ³)	Do not award 25.02 (cm ³) for mean of all three	
		titres	

Question	Answer	Additional Guidance	Mark
Number			
5(c)(iii)		Example of calculation	(1)
	moles of hydrochloric acid in the titration	$24.90 \times 0.10 \div 1000 = 2.49 \times 10^{-3} / 0.00249 $ (mol) TE on 5(c)(ii) Ignore SF except 1SF	

Question Number	Answer		Additional Guidance	Mark
5(c)(iv)		Example	e of calculation	(2)
	 moles of M₂CO₃ in 25 cm³ of solution 	(1) 2.49 × 1 TE on 5($0^{-3} \div 2 = 1.245 \times 10^{-3} / 0.001245$ (mol) (c)(iii)	
	 moles of M₂CO₃ in 250 cm³ of solution 	TE on M	$10^{-3} \times 10 = 1.245 \times 10^{-2} / 0.01245 $ (mol) 11 SF except 1SF	

Question	Answer	Additional Guidance	Mark
Number			
5(c)(v)		Example of calculation	(1)
	 molar mass of M₂CO₃.10H₂O 	$3.56 \div 1.245 \times 10^{-2} = 285.9 / 286 \text{ (g mol}^{-(1)})$ TE on 5(c)(iv) Allow g for g mol $^{-(1)}$ Ignore SF except 1SF	

Question Number	Answer		Additional Guidance	Mark
5(c)(vi)	 subtraction of mass of CO₃ and 10H₂O from molar mass 	(1)	Example of calculation 286 – 180 – 48 – 12 (= 46) TE on 5(c)(v)	(2)
	identification of metal, M	(1)	Dependent on M1 (or attempt at subtraction of carbonate and water) = $46 \div 2 = 23$ (g mol ⁻⁽¹⁾) and Na	
			Allow g for g mol ⁻⁽¹⁾ Just Na with no working scores 0	
			Metal must be identified as the Group 1 element with molar mass closest to the calculated value	

Question Number	Answer		Additional Guidance	Mark
5(d)	A description that makes reference to the following points:		Penalise use of indicator once only Penalise use of drying agent in M1 and M3 once only	(3)
	 evaporate/heat and to crystallisation/saturation (point) leave to (stand and) crystallise / cool 	(1) (1)	Allow evaporate some/half/most of the water Allow heat until solid starts to form Allow heat to concentrate the solution Do not award evaporate to dryness Do not award heat to constant mass Do not award remove some water with drying agent	
	(decant the supernatant liquid and) dry crystals between filter paper	(1)	Allow dry between tissue / in a desiccator/oven	
			Ignore reference to washing of crystals prior to drying Do not award dry crystals with drying agent	

(Total for Question 5 = 14 marks)

TOTAL FOR PAPER = 50 MARKS