Please check the examination details below Candidate surname	Other names
Pearson Edexcel International Advanced Level	Candidate Number
Friday 18 Janua	ry 2019
Afternoon (Time: 1 hour 15 minutes)	Paper Reference WCH03/01
Chemistry Advanced Unit 3: Chemistry Laborator	ry Skills I
Candidates must have: Scientific calc	ulator Total Marks

Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
 - there may be more space than you need.

Information

- The total mark for this paper is 50.
- The marks for **each** question are shown in brackets
 - use this as a guide as to how much time to spend on each question.
- You will be assessed on your ability to organise and present information, ideas, descriptions and arguments clearly and logically, including your use of grammar, punctuation and spelling.
- A Periodic Table is printed on the back cover of this paper.

Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ▶







Answer ALL the questions. Write your answers in the spaces provided.

- 1 A white solid A contains one cation and one anion.
 - (a) A small amount of solid **A** was placed in a test tube and aqueous sodium hydroxide added. The mixture was warmed gently.

Complete the inference column in the table.

(2)

Observation	Inference
A pungent smelling gas was evolved that turned damp red litmus paper blue	The gas formed is
	The formula of the cation in A is

(b) (i) An aqueous solution of **A** was placed in a test tube and acidified with dilute nitric acid. A few drops of silver nitrate solution were added. Complete the inference column in the table.

(1)

Observation	Inference
Cream precipitate formed	The precipitate is

(ii) Write the **ionic** equation, including state symbols, for the formation of the cream precipitate in (b)(i).

(2)



(iii) Describe how you would confirm the identity of formed in (b)(i).	the anion in the cream precipitate
ioimed iii (b)(i).	(2)
	(Total for Question 1 = 7 marks)

2 (a) A student was provided with aqueous solutions of four compounds:

barium nitrate

hydrochloric acid

sodium carbonate

sulfuric acid

Four bottles, labelled **B**, **C**, **D** and **E**, each contained one of the solutions. The student mixed pairs of the solutions to determine which solution was in each bottle.

The results are shown.

Solutions mixed	Observations
B and C	Effervescence with bubbles of a colourless gas given off
B and D	No visible change
B and E	A white precipitate formed which did not dissolve on the addition of dilute nitric acid
C and D	Effervescence with bubbles of a colourless gas given off
C and E	A white precipitate formed which dissolved with effervescence on the addition of dilute nitric acid
D and E	No visible change

Use the observations in the table to deduce the identity of the compound in each bottle. Identify each compound by name or formula.

I	3																	

(3)

	Describe how you would carry out a flame test.	
	Describe now you would carry out a name test.	(3)
(ii)	State the flame colours produced by barium nitrate and sodium carbonate.	
	Barium nitrate	
	Sodium carbonate	(2)
	(Total for Question 2 = 8 r	narks)



3 Chlorine-based bleaches contain sodium chlorate(I), NaClO, as the active ingredient. The concentration of NaClO in bleach was determined by a titration method using sodium thiosulfate.

Sodium chlorate(I) reacted with potassium iodide in acidic solution to produce iodine.

$$ClO^{-} + 2I^{-} + 2H^{+} \rightarrow I_{2} + Cl^{-} + H_{2}O$$

The iodine was then titrated with sodium thiosulfate.

$$2S_2O_3^{2-} + I_2 \rightarrow 2I^- + S_4O_6^{2-}$$

Procedure

- 1. A burette was filled with 0.0600 mol dm⁻³ sodium thiosulfate solution.
- 2. 10.0 cm³ of bleach was pipetted into a 250.0 cm³ volumetric flask and excess potassium iodide and sulfuric acid were added to release iodine. The volume was made up to the mark with distilled water.
- 3. 25.0 cm³ of this solution was pipetted into a conical flask and titrated with the sodium thiosulfate solution using a suitable indicator.
- (a) State the indicator used and give the colour change at the end-point.

(2)

Indicator	Colour change at the end-point							
	From	to						

(b) (i) Complete the table of results.

(1)

Number of titration	1	2	3	4
Burette reading (final) / cm ³	23.65	46.45	24.40	47.10
Burette reading (start) / cm ³	0.00	23.65	1.20	24.40
Titre / cm ³				

(ii) State with a reason which results should be used to calculate the mean titre value.

(2)

(iii) Calculate the mean titre.

(1)

(iv) Calculate the number of moles of sodium thiosulfate in this mean titre.

(1)

(v) Calculate the number of moles of iodine in 25.0 cm³ of the diluted solution.

(1)

(vi) Calculate the number of moles of sodium chlorate(I) in the 250.0 cm³ volumetric flask.

(1)

(vii) Calculate the concentration of sodium chlorate(I) in the **undiluted** bleach in mol dm⁻³.

(1)



(0	The $0.0600\mathrm{moldm^{-3}}$ sodium thiosulfate solution used in this titration is known as a standard solution.	
	Describe the steps you would take to prepare this standard solution as accurately as possible. You are supplied with the appropriate mass of sodium thiosulfate and the usual laboratory glassware, including a volumetric flask.	
	No calculations are required.	
	The calculations are required.	(3)
		(-)
	(Total for Question 3 = 13 ma	rke)
		I N3/

BLANK PAGE




4 Hydrogen peroxide, H₂O₂, decomposes according to the equation

$$2H_2O_2 \rightarrow 2H_2O + O_2$$

The rate of decomposition is increased by a catalyst.

A student tested three metal oxides to determine which was the best catalyst. The oxides were manganese(IV) oxide, iron(III) oxide and lead(IV) oxide. They are all solids.

The student used the following apparatus and experimental procedure.



Procedure


- 1. Hydrogen peroxide solution was poured into the conical flask.
- 2. Solid manganese(IV) oxide was added.
- 3. The bung was quickly replaced to connect the gas syringe to the conical flask.
- 4. The procedure was repeated using iron(III) oxide and lead(IV) oxide.

when using this procedure.	(3)
(b) State the measurements the student should make to determine which	is the
best catalyst.	(2)
(c) The student thought that some of the gas assemed from the conical fla	sk boforo
(c) The student thought that some of the gas escaped from the conical fla the bung had been replaced.	sk belole
Suggest how this experiment could be modified to prevent this loss.	(5)
	(1)



(d) Another student thought that some of the oxygen produced may have come from the decomposition of the metal oxide.	
Suggest how this idea could be tested.	(2)
(Total for Question 4 = 8 mai	'ks)

5 Some organic reactions are shown.



(a) Reaction 1 and Reaction 2 use the same reagent but require different conditions. Identify the reagent and give the conditions needed for Reaction 1.

(2)

(b) (i) Give a chemical test and its positive result to show the presence of the double bond in $CH_3CH = CH_2$.

(2)

(ii) Give the structure of the organic product of the test in (b)(i).

(1)



(c)	A student added phosphorus(V) chloride, PCl ₅ , to the product of Reaction 2 ,
	CH₃CH₂CH₂OH. Hydrogen chloride was formed.

(i) S	State the	observation	the student	would be e	expected to	make.
-------	-----------	-------------	-------------	------------	-------------	-------

(1)

(ii) Complete the table to show the hazard and the appropriate safety precaution for each chemical.

Do not include the wearing of eye protection and a laboratory coat.

(3)

Chemical	Hazard	Safety precaution
PCl₅		
CH ₃ CH ₂ CH ₂ OH		
HCl		

- (d) In Reaction **3**, CH₃CH₂CH₂OH is oxidised to CH₃CH₂CHO using aqueous potassium dichromate(VI) acidified with sulfuric acid.
 - (i) State the colour **change** that occurs during this oxidation reaction.

(1)



ps://t.me/joinchat/wwc3WbVZ6MtkYWU0	www.igexams.com
(ii) Draw a labelled diagram of the apparatus you would u	se to carry out
Reaction 3 and collect the product.	•
	(3)
(iii) Explain how infrared spectroscopy could be used to co	
CH ₃ CH ₂ CH ₂ OH has been oxidised to CH ₃ CH ₂ CHO in Rea	ction 3 .
You are not expected to give specific wavenumbers.	(1)
	(1)

(Total for Question 5 = 14 marks)

TOTAL FOR PAPER = 50 MARKS



Er erbium 68

Tb terbium 9

9

59

64

63

62

99

69

[257]

103

102

mendelevium 101

einsteinium 99

californium 98

PW

[253] Fm fermium 100

[254] 49

[251] Cf

[245] BK berketium 97

E supplemental

americium Am 95

Np neptunium 93

uranium

otactinium

92

91

94

Pu Pu plutonium

[237] 9

[231]

Ce certum 58 232 Th Th thorium 90

[247]

nobelium

+	2
2	1)
8	
0	با
L	_
4	5
0	U
(5
۴	O
2	ر
3	2
.5	_
0	U
0	U
Ļ	

0 (8)

1

9

2

m

				_					_				_								_				rs						
(18)	4.0	He	netium,	7	20.2	Ne	neon	10	39.9	Ar	argon	18	83.8	ᅶ	krypton	36	131.3	Xe	xenon	25	[222]	R	radon	98	1	pa					
•			(47)		19.0	ı	fluorine	6	35.5	บ	chlorine	17	6.62	В	bromine	35	126.9	-	iodine	53	[210]	At	astatine	85		een report			175	2	lutetium
			(347)	(6)	16.0	0	oxygen	80	32.1	S	sulfur	16	79.0	Se	selenium	34	127.6	Te	tellurium	52	[509]	8	polonium	84	0.000	116 have b	ticated		173	χ	ytterbium
			1461	6	14.0	z	nitrogen	7	31.0	۵	phosphorus	15	74.9	As	arsenic	33	121.8	Sb	antimony	51	209.0	Bi	bismuth	83	ASSESSED FOR	mbers 112-	ully authen		169	Ē	thulium
			(117)	(+1)	12.0	U	carbon	9	28.1	Si	_	14	72.6	ge	germanium	32	118.7	Sn	tin	20	207.2	Ъ	lead	82	8	atomic nur	but not fi		167	ū	erbium
			1431	(c)	10.8	В	poron	2	27.0	AI	aluminium	13	69.7	Ga	gallium	31	114.8	'n	indium	49	204.4	F	thallium	81	2000	nents with			165	e H	ž
				3								(12)	65.4	Zn	zinc	30	112.4	В	cadmium	48	200.6	Hg	mercury	80	was	Elen			163	۵	dysprosium
												(11)	63.5	D	copper	53	107.9	Ag	silver	47	197.0	Αn	plog	79	[272]	Rg	roentgenium	111	159	4	E
												(10)	58.7	ž	nickel	28	106.4	Pd	palladium	46	195.1	£	platinum	78	[271]	Os	darmstadtium	110	157	PS	gadolinium
											į	(6)	58.9	ပိ	cobalt	27	102.9	뫈	rhodium	45	192.2	<u>_</u>	iridium	77	[392]	¥	meltnerium	109	152	Eu	E
•	?:	hydrogen	`-								9	(8)	55.8	Fe	iron	56	101.1	Ru	ruthenium	44	190.2	o	osmium	9/	[277]	¥	F	108	150	Sm	samarium
											į	(2)	54.9	Wn	manganese	25	[86]	2	technetium	43	186.2	Re	rhenium	75	400			107	[147]	Pm	praseodymium promethium samarium
					mass	lod		umber				(9)	52.0	Ն	chromium	24	626	Wo	molybdenum	42	183.8	>	tungsten	74	[592]	Sg	seaborgium	106	144	P	neodymium
			Key ve atomic r mic symb name (proton) nu				we atomic mic syml name (proton) n					(2)	50.9	>	vanadium	23	92.9	å	niobium	41	180.9	Тa	tantalum	73	_		E	105	141	P	praseodymium
					relati	ato		atomic			3	(4)	47.9	F	titanium	22	91.2	Zr	zirconium	40	178.5	Ŧ	hafnium	72	[261]	¥	nutherfordium	104	140	ő	-
											ļ	(3)	45.0	Sc	scandium	21	88.9	>	yttrium	39	138.9	La*	Lanthanum	22	[227]	Ac*	E	68		52	
			10/	(7)	0.6	Be	beryllium	4	24.3	Wa	magnesium	12	40.1	Ca	calcium	20	87.6	S	strontium	38	137.3	Ba	F	99	[526]	Ra	radium	88		anide serie	* Actinide series
			140	3	6.9	5	lithium	٣	23.0	Na	0	11	39.1	¥	potassium	19	85.5	&	rubidium	37	132.9	క	caesium	55	[223]	占	francium	87		* Lanth	* Actinic
				H hydrogen 1.0	1.0 H hydrogen 1 (13) (14) (15) (16) (17)	(2) Key (13) (14) (15) (16) (17) (17) (19.0 19.0	(2) Key (13) (14) (15) (16) (17) (19.0 P. Be atomic symbol	(2) Key (13) (14) (15) (16) (17) (17) (19	(2) Key 1.0 H hydrogen 1.0 14.0 15.0 17.0 19.0	(2) Key 1.0 H hydrogen 1.0 H hydrogen 1.0 1.	(2) Key 1.0 Hydrogen 1.0 Telative atomic mass atomic (proton) number 24.3 Al Si P S CI May Si P S CI Hydrogen 1.0 Hydrogen	(2) Key 1.0 H hydrogen 1.0 9.0 relative atomic mass atomic (proton) number 4 atomic (proton) number Am atomic (proton) number Am atomic (proton) number Am Si P S CI Am Si Suffur chlorine allowinishm silicon phosphorus suffur chlorine suffur chlorine atomic prospersium silicon phosphorus suffur chlorine suffur chlor	(2) Key 1.0 Hydrogen 1.0 Hydrog	1.0 Hydrogen 1.0	(2) Key Telative atomic mass atomic (proton) number A 40.1 (5.0 (6) (7) (8) (9) (10) (11) (12) (12) (14) (15) (16) (17) (17) (18) (19) (19) (19) (19) (19) (19) (19) (19	1.0 Holydrogen Holydrogen 1.0 Holydrogen 1.0 Holydrogen Holydrogen Holydrogen 1.0 Holydrogen Holydro	1.0 Hydrogen 1.0	1.0 Hydrogen 1.0	1.0 Hydrogen Hydrogen 1.0 Hydrogen 1.0 Hydrogen 1.0 Hydrogen Hydrogen	1.0 Hydrogen H	1.0 H H H H H H H H H	1.0 H H H H H H H H H	1.0 H H H H H H H H H	2,2 2,2	1.0 1.0	1.0 Hydrogen Hydrogen 1.0 Hydrogen 1.0 Hydrogen 1.0 Hydrogen Hydrogen 1.0 Hydrogen	2) 2) 2) 2) 2) 2) 2) 2)	20 1.00 1.	1.0 Holdrogen 1.0 Hold	1.0 February Feb	(2) 8.00 8.00 8.00 8.00 8.00 8.00 8.00 8.0

P 5 1 7 0 7 A 0 1 6 1 6