



Mark Scheme (Results)

June 2014

GCE Chemistry (6CH04/01)
General Principles of Chemistry I

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at www.edexcel.com.

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

www.edexcel.com/contactus

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at:

www.pearson.com/uk

Summer 2014

Publications Code UA038324

All the material in this publication is copyright

© Pearson Education Ltd 2014

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Mark schemes will indicate within the table where, and which strands of QWC, are being assessed. The strands are as follows:
 - i) ensure that text is legible and that spelling, punctuation and grammar are accurate so that meaning is clear
 - ii) select and use a form and style of writing appropriate to purpose and to complex subject matter
 - iii) organise information clearly and coherently, using specialist vocabulary when appropriate

Using the Mark Scheme

Examiners should look for qualities to reward rather than faults to penalise. This does NOT mean giving credit for incorrect or inadequate answers, but it does mean allowing candidates to be rewarded for answers showing correct application of principles and knowledge. Examiners should therefore read carefully and consider every response: even if it is not what is expected it may be worthy of credit.

The mark scheme gives examiners:

- an idea of the types of response expected
- how individual marks are to be awarded
- the total mark for each question
- examples of responses that should NOT receive credit.

/ means that the responses are alternatives and either answer should receive full credit.

() means that a phrase/word is not essential for the award of the mark, but helps the examiner to get the sense of the expected answer.

Phrases/words in **bold** indicate that the meaning of the phrase or the actual word is **essential** to the answer.

ecf/TE/cq (error carried forward) means that a wrong answer given in an earlier part of a question is used correctly in answer to a later part of the same question.

Candidates must make their meaning clear to the examiner to gain the mark. Make sure that the answer makes sense. Do not give credit for correct words/phrases which are put together in a meaningless manner. Answers must be in the correct context.

Quality of Written Communication

Questions which involve the writing of continuous prose will expect candidates to:

- write legibly, with accurate use of spelling, grammar and punctuation in order to make the meaning clear
- select and use a form and style of writing appropriate to purpose and to complex subject matter
- organise information clearly and coherently, using specialist vocabulary when appropriate.

Full marks will be awarded if the candidate has demonstrated the above abilities.

Questions where QWC is likely to be particularly important are indicated (QWC) in the mark scheme, but this does not preclude others.

Section A (multiple choice)

Question Number	Correct Answer	Reject	Mark
1 (a)	B		1

Question Number	Correct Answer	Reject	Mark
1 (b)	C		1

Question Number	Correct Answer	Reject	Mark
1 (c)	A		1

Question Number	Correct Answer	Reject	Mark
1 (d)	D		1

Question Number	Correct Answer	Reject	Mark
2 (a)	B		1

Question Number	Correct Answer	Reject	Mark
2 (b)	D		1

Question Number	Correct Answer	Reject	Mark
2 (c)	D		1

Question Number	Correct Answer	Reject	Mark
2 (d)	C		1

Question Number	Correct Answer	Reject	Mark
3	D		1

Question Number	Correct Answer	Reject	Mark
4 (a)	B		1

Question Number	Correct Answer	Reject	Mark
4 (b)	B		1

Question Number	Correct Answer	Reject	Mark
4 (c)	C		1

Question Number	Correct Answer	Reject	Mark
5	B		1

6 (a)	B		1
--------------	---	--	----------

Question Number	Correct Answer	Reject	Mark
6 (b)	C		1

Question Number	Correct Answer	Reject	Mark
6 (c)	D		1

Question Number	Correct Answer	Reject	Mark
6 (d)	A		1

Question Number	Correct Answer	Reject	Mark
7 (a)	A		1

Question Number	Correct Answer	Reject	Mark
7 (b)	C		1

Question Number	Correct Answer	Reject	Mark
8	A		1

Section B

Question Number	Acceptable Answers	Reject	Mark
9 (a) (i)	+89.6 – [+32.7 + 165] (1)		2
	= –108.1 J mol ⁻¹ K ⁻¹ / J K ⁻¹ mol ⁻¹		
	Value, sign and units (1)		
	Ignore SF except one		
	Internal TE for recognisable numbers allowed, for example:		
	$\Delta H^\ominus_{\text{at}}$ magnesium chloride (147.7 → -223.1)		
Halving S^\ominus [Cl ₂] (82.5 → -25.6)			
Correct answer with no working (2)			
+ /no sign 108.1 J mol ⁻¹ K ⁻¹ / J K ⁻¹ mol ⁻¹ (1)			

Question Number	Acceptable Answers	Reject	Mark
9 (a) (ii)	<p>(The sign is negative because)</p> <p>Any two from:</p> <ul style="list-style-type: none"> • (A solid and) a gas reacting to form a solid. <p>OR</p> <p>(Entropy decreases because) a gas reacting to form a solid.</p> <ul style="list-style-type: none"> • There are fewer ways of arranging particles in a solid than a gas or vice-versa. <p>OR</p> <p>Decrease in disorder as solid more ordered than gas or vice versa</p> <ul style="list-style-type: none"> • Two mol(es) of reactant forming one mole of product. (Ignore two molecules form one molecule) <p>OR</p> <p>Number of mol(es)/molecules decreases</p> <p>OR</p> <p>Fewer/less mol(es) of products than reactants</p>	<p>Energy...</p> <p>'(Positive) Answer is as expected...' (0)</p>	2

Question Number	Correct Answer	Reject	Mark
9 (b)	$\Delta S^{\circ}_{\text{total}} = \Delta S^{\circ}_{\text{surroundings}} + \Delta S^{\circ}_{\text{system}}$ <p>OR</p> $= +2152 + (-108.1)$ $= (+)2043.9$ <p>Value 2043.9 / 2044 (1)</p> $= (+)2040 \text{ (J mol}^{-1} \text{ K}^{-1}\text{)}$ <p>3SF</p> <p>This mark conditional on correct value or correct TE value from (a)(i) (1)</p> <p>Accept TE from (a)(i), for example,</p> $-223.1 \rightarrow +1928.9 \rightarrow +1930$ $-25.6 \rightarrow +2126.4 \rightarrow +2130$ <p>Correct answer (2040, etc) with or without working scores 2</p>		2


Question Number	Correct Answer1	Reject	Mark
9 (c)	$\Delta S_{\text{surroundings}}^{\ominus} = - \frac{\Delta H^{\ominus}}{298}$ $\Delta H^{\ominus} = - \Delta S_{\text{surroundings}}^{\ominus} \times 298$ <p>OR</p> $\Delta H^{\ominus} = -2152 \times 298 \quad (1)$ $= -641.296$ $= -641.3 \text{ (kJ mol}^{-1}\text{)} \quad (1)$ <p>ALLOW</p> $= -641.3 \times 10^3 \text{ J mol}^{-1}$ <p>Note</p> <ol style="list-style-type: none"> -640.1338 = -640.1 (if 2040/answer to part (b) used to recalculate entropy change of surroundings first.) (2) $\Delta H^{\ominus} = +641.3 \text{ (kJ mol}^{-1}\text{)} \quad (1)$ $\Delta H^{\ominus} = - \frac{\Delta S_{\text{surroundings}}^{\ominus}}{298} \quad (0)$ <p>Ignore SF except one</p>		2

Question Number	Correct Answer	Reject	Mark
9 (d) (i)	$50 \times 4.2 \times 22.5$ = 4725 (J) Ignore sign ALLOW 4.725 kJ Ignore SF except one		1

Question Number	Correct Answer	Reject	Mark
9 (d) (ii)	There are two legitimate answers to this part. If both methods have been used, you must send the item to review under mark scheme $(-)\text{4725} \div 0.0300$ = $-157.5 \text{ (kJ mol}^{-1}\text{)}$ / $-157500 \text{ J mol}^{-1}$ OR $(-)\text{4725} \div 0.0500$ = $-94.5 \text{ (kJ mol}^{-1}\text{)}$ / $-94500 \text{ J mol}^{-1}$ ALLOW TE answer (d)(i) $\div 0.0300/0.0500$ Ignore SF except one Value (1) Sign (1) The mark for the negative sign is awarded for their calculation even if value is wrong, providing any energy divided by moles or energy multiplied by 1/number of moles calculation has been done.		2

Question Number	Correct Answer	Reject	Mark
9 (d) (iii)	<p>There are two correct answers:</p> <p>Using 0.03 gives the answer of $-381.75 \text{ kJ mol}^{-1}$</p> <p>Using 0.05 gives the answer of $-350.25 \text{ kJ mol}^{-1}$</p> <p>Both these answers score full marks with or without correct working.</p> <p>First mark</p> <p>Appreciation of Hess's Law either in words, numbers, symbols or on the diagram</p> <p>For example,</p> $\Delta H_{\text{solution}} + \text{Lattice energy}$ $= \Delta H_{\text{hydration}} \text{Mg}^{2+} + (2)\Delta H_{\text{hydration}} \text{Cl}^{-}$ <p style="text-align: right;">(1)</p> <p>Second mark</p> $2 \Delta H_{\text{hydration}} \text{Cl}^{-} = -2526 - 157.5 - (-1920) = -763.5$ <p>OR</p> $2 \Delta H_{\text{hydration}} \text{Cl}^{-} = -2526 - 94.5 - (-1920) = -700.5$ <p>ALLOW</p> <p>Any number or group of numbers minus (-1920) (1)</p> <p>Third mark</p> $\Delta H_{\text{hydration}} \text{Cl}^{-} = -381.75 \text{ (kJ mol}^{-1}\text{)}$ <p>OR</p> $\Delta H_{\text{hydration}} \text{Cl}^{-} = -350.25 \text{ (kJ mol}^{-1}\text{)}$ <p>Any number, wherever it has come from,</p>		3

	divided by two can score this mark, provided that the sign is consistent.	(1)		
	Ignore SF except one			
	Use of lattice energy – 2326 gives –281.75/–250.25 scores	(2)		
	ALLOW			
	TE from (d)(ii)			

Question Number	Correct Answer	Reject	Mark
9 (d)(iv)	 <p style="text-align: center;">OR</p> <ul style="list-style-type: none"> • One/several water molecule(s) all correctly orientated. • $H^{\delta+}$/ hydrogen (one or two hydrogens from each water molecule) towards chloride ion • with negative charge either on chlorine or on the whole hydrated ion. <p>ALLOW</p> <ul style="list-style-type: none"> • A minus sign with a ring around it for the Cl^- • Bonds shown by lines/broken lines/dotted lines/wedges 	<p style="text-align: center;">$Cl^- \cdot H_2O$</p> <p style="text-align: center;">$H^{\delta-} / H^+ / H^-$</p> <p style="text-align: center;">$Cl^{\delta-} / Cl$ (with no charge)</p>	1

Question Number	Correct Answer	Reject	Mark
9(d)(v)	<p>Both marks may be awarded in either part.</p> <p>First mark</p> <p>(Temperature increases) because the reaction/process/dissolving/hydration of ions is exothermic.</p> <p>OR</p> <p>Strong(er) forces between the δ^+ H and Cl^-</p> <p>OR</p> <p>Strong(er) forces between the δ^- O and Mg^{2+}</p> <p>OR</p> <p>Strong(er) ion-dipole forces</p> <p>OR</p> <p>Formation of bonds releases energy</p> <p>OR</p> <p>Strong(er) bonds formed</p> <p>OR</p> <p>Enthalpy of hydration is greater than lattice energy (1)</p> <p>Second mark</p> <p>(Volume decreases so) shorter bonds between ion and water molecules</p> <p>ALLOW</p> <p>Water molecules more tightly arranged/pack better/occupy less space</p> <p>OR</p> <p>Water molecules more ordered/ clustered (around the ions). (1)</p>	<p>The breaking of the lattice is exothermic.</p> <p>Ions more tightly arranged</p> <p>Ions more ordered</p>	2

Total for Question 9 = 17 marks

Question Number	Correct Answer	Reject	Mark
10(a)(i)	<p>Sodium/potassium dichromate ((VI)) and (Dilute/concentrated) sulfuric acid</p> <p>OR</p> <p>correct formulae / H^+ and $\text{Cr}_2\text{O}_7^{2-}$</p> <p>ALLOW</p> <p>$\text{H}^+$ and $\text{Cr}_2\text{O}_7^{2-}$/acidified dichromate((VI)) (1)</p> <p>Reflux/distil</p> <p>Ignore 'heat', 'warm', and 'boil' alone.</p> <p>ALLOW</p> <p>Just 'under reflux'</p> <p>Just 'under distillation' (1)</p> <p>Second mark depends on mention of dichromate/$\text{Cr}_2\text{O}_7^{2-}$ in first part</p> <p>OR</p> <p>KMnO_4 and acid with heat (1)</p>	Hydrochloric acid	2

Question Number	Correct Answer	Reject	Mark
10 (a)(ii)	<p>Carbonyl group – addition of 2,4-dinitrophenylhydrazine / 2,4-DNP(H) / Brady's reagent (1)</p> <p>to give yellow/orange/red precipitate/ppt/ppte/solid/crystals</p> <p>ALLOW</p> <p>recognisable spelling e.g., percepitate (1)</p> <p>CH₃C=O reaction with iodine in alkali/NaOH/KOH/OH⁻</p> <p>ALLOW</p> <p>Iodoform/tri-iodomethane/haloform</p> <p>AND</p> <p>reaction/test (1)</p> <p>to form (pale) yellow / cloudy precipitate/solid/crystals (1)</p> <p>Ignore references to smell</p> <p>Ignore heat in either part</p> <p>Note</p> <ul style="list-style-type: none"> In both cases result mark depends on test being recognisably correct even if it did not score a mark <p>Examples:</p> <p>DNP gives yellow ppt</p> <p>Iodine test gives yellow ppt</p> <ul style="list-style-type: none"> Tests for aldehydes with correct results, no marks 	<p>2-DNP/4DNP</p> <p>Just DNP</p> <p>Brick red ppt</p>	4

Question Number	Correct Answer	Reject	Mark
10 (b) (i)	<p>Arrow (from carbon) of CN^- to carbon of $\text{C}=\text{O}$</p> <p>AND</p> <p>Arrow from part of $\text{C}=\text{O}$ double bond to oxygen</p> <p>ALLOW</p> <p>Two steps via a charged canonical form (1)</p> <p>Intermediate anion with $\text{C}-\text{CN}$ bond. (1)</p> <p>Arrow from resulting O^- to hydrogen of $\text{HCN}/\text{H}^+/\text{H}_2\text{O}$ (1)</p> <p>Note</p> <p>Arrow directions must be correct to score each mark Penalise half-headed arrows each time in both parts ALLOW skeletal formulae.</p>	<p>CN without negative charge</p> <p>...C-NC bond</p>	3


Question Number	Correct Answer	Reject	Mark
10 (b) (ii)	<p>Forms a racemic mixture (1)</p> <p>Because bonds around C=O are planar</p> <p>OR</p> <p>Carbonyl group/reaction site is (trigonal) planar</p> <p>OR</p> <p>Bonds around carbonyl carbon are planar (1)</p> <p>Cyanide can attack from either side / above or below (1)</p>	<p>Butanone/molecule/it is planar</p> <p>C=O is planar</p> <p>Carbonyl bond is planar</p> <p>Intermediate is planar</p>	3

Question Number	Correct Answer	Reject	Mark
10 (c) (i)	<p>(Acid) hydrolysis</p> <p>OR</p> <p>Alkaline hydrolysis followed by acidification</p>	Hydration	1

Question Number	Correct Answer	Reject	Mark
10 (c) (ii)	<p>The O-H absorptions for alcohol and carboxylic acid overlap.</p> <p>OR</p> <p>OH absorption for an acid is very broad</p> <p>OR</p> <p>Quote data booklet values which must show some overlap, to include 3300 to 3200.</p> <p>ALLOW</p> <p>OH absorptions similar/the same.</p>	<p>Just 'both have OH groups'</p> <p>Just 'two OH groups present'</p>	1

Question Number	Correct Answer	Reject	Mark
10 (c) (iii)	<p>(Chemical shift δ) 2.0 - 4.0 (ppm) / any value within this range</p> <p>ALLOW</p> <p>Correct number followed by δ, eg 3δ</p>		1

Question Number	Correct Answer	Reject	Mark
10 (c) (iv)	<p>There is no hydrogen atom/proton on the adjacent/neighbouring carbon atom</p> <p>ALLOW</p> <p>No adjacent/neighbouring hydrogens/protons</p>		1

Question Number	Correct Answer	Reject	Mark
10 (d)	 <p data-bbox="397 850 1128 892">Ester linkage (1)</p> <p data-bbox="397 913 1128 955">Rest of molecule (1)</p> <p data-bbox="397 1018 495 1050">ALLOW</p> <p data-bbox="397 1081 933 1113">Attached chains as structural formulae</p> <p data-bbox="397 1144 990 1176">Ignore n or other numbers outside bracket</p>		2


Total for Question 10 = 18 marks

Question Number	Correct Answer	Reject	Mark
11(a)	$\text{S}_2\text{O}_8^{2-} + 2\text{I}^- \rightarrow 2\text{SO}_4^{2-} + \text{I}_2$ <p>ALLOW multiples</p> <p>Ignore state symbols even if incorrect</p>		1

Question Number	Correct Answer	Reject	Mark
11 (b) (i)	<p>Blue/black /blue-black</p> <p>OR</p> <p>Colourless to blue-black/ blue/black</p>	Purple	1

Question Number	Correct Answer	Reject	Mark
11 (b) (ii)	<p>The mixture would change colour/ go blue/black /blue-black immediately/straight away</p> <p>ALLOW</p> <p>...too quick(ly)/too early</p> <p>...quicker</p> <p>...no time delay</p>		1

Question Number	Correct Answer	Reject	Mark
11 (b) (iii)	<p>(As quickly as iodide reacts to form iodine it is) reduced/turned back to iodide by the thiosulfate ions</p> <p>ALLOW</p> <p>Persulfate reacts with thiosulfate first.</p> <p>OR</p> <p>Iodine reacts with thiosulfate.</p>		1

Question Number	Correct Answer	Reject	Mark
<p>11 (c) (i)</p>	 <p>First mark Correct graph of rate v concentration, with axes correct and values increasing on both axes labelled with quantity and units Note Units may be given in brackets with no slash. s/time meaning s divided by time is fine. (1)</p> <p>Second mark Sensible scales to use at least half the graph paper but allow graphs starting at the origin and points cover two by two big squares. Linear scales All points reasonably correct with straight line drawn (1) Second mark depends on correct graph of rate v concentration, but not other detail of first mark</p>		2

Question Number	Correct Answer	Reject	Mark
11 (c) (ii)	<p>First order</p> <p>This mark is independent of the graph drawn (1)</p> <p>Because the graph is a straight line (through the origin)/ rate is proportional to $[S_2O_8^{2-}]$</p> <p>OR</p> <p>As concentration increases by (factor of) 2 rate increases by 2 (or any other numbers, including 'x')</p> <p>OR</p> <p>Rate increases linearly (with concentration)</p> <p>OR</p> <p>Gradient of line is constant (1)</p> <p>Second mark depends on first order</p>	Just 'as concentration increases rate increases'	2

Question Number	Correct Answer	Reject	Mark
11 (c) (iii)	<p>Rate = $k[S_2O_8^{2-}][I^-]$ (1)</p> <p>TE from (c)(ii)</p> <p>Units - $dm^3 mol^{-1} s^{-1}$</p> <p>ALLOW</p> <p>Internal TE from rate equation (1)</p> <p>Units in any order</p>	Incorrect formulae	2

Question Number	Correct Answer	Reject	Mark
11 (d)(i)	<p>Method 1</p> <p>First mark</p> <p style="padding-left: 40px;">Gradient = $- E_a/R$</p> <p>OR</p> <p style="padding-left: 40px;">$E_a = - R \times \text{gradient}$ (1)</p> <p>Second mark</p> <p style="padding-left: 40px;">(Gradient =) $\frac{-3.0 - (-3.69)}{(3.30 - 3.41) \times 10^{-3}}$</p> <p>OR</p> <p style="padding-left: 80px;">= -6272.7 (K)</p> <p>Please award this mark if -6272.7 is seen anywhere! (1)</p> <p>Method 2</p> <p>First mark</p> <p>Setting up two simultaneous equations (1)</p> <p>Second mark</p> <p>Subtracting one equation from the other or other correct methods of solution (1)</p> <p>Third mark (applies to both methods)</p> <p style="padding-left: 40px;">$(E_a) = +52126 \text{ J mol}^{-1}$ $\quad \quad \quad / +52.1(26) \text{ kJ mol}^{-1}$</p> <p>Note: TE can only be given if either method 1 or method 2 has been clearly carried out.</p> <p>Positive sign given</p> <p>OR</p> <p>Two negative signs clearly cancel in method and no sign given (1)</p> <p>Correct answer with or without working, with sign and units (3)</p> <p>Ignore SF unless only one</p>	<p>Negative sign</p>	<p>3</p>

Question Number	Correct Answer	Reject	Mark
11 (d)(ii)	Either Take readings at different temperatures OR Repeat at the same two temperatures ALLOW Just 'repeat the experiment'		1

Total for Question 11 = 14 marks

Total for Section B = 49 marks

Section C

Question Number	Correct Answer	Reject	Mark
12(a)(i)	<p>Mass of ethanoic acid = 0.04×60.1 = (2.404 g) (1)</p> <p>Volume of ethanoic acid = $2.404 \div 1.049$ = 2.2917 = 2.3 (cm³) (1)</p> <p>Correct answer with no working (2)</p> <p>Ignore SF except only one</p> <p>ALLOW</p> <p>60.0 for molar mass which gives mass 2.4 and volume 2.288 = 2.3 cm³ (2)</p> <p>OR</p> <p>First step $1.049 \div 60/60.1$ to find number of moles in 1 cm³ = 0.017 (1)</p> <p>Then volume = $0.04 \div 0.017$ = 2.3529 (cm³)</p> <p>But note, if whole calculation done on calculator, 60 gives 2.2879 and 61 gives 2.2917. (1)</p> <p>If units given, they must be correct, but penalise wrong units only once here.</p>		2

Question Number	Correct Answer	Reject	Mark
12(a)(ii)	<p>Syringe</p> <p>ALLOW Burette</p> <p>Graduated/adjustable pipette</p>	<p>Gas syringe</p> <p>Biuret</p> <p>Just 'pipette'</p>	1

Question Number	Correct Answer	Reject	Mark
12 (a)(iii)	To prevent... evaporation/vapour escaping water vapour entering OR To maintain a closed system OR To maintain a closed environment ALLOW To prevent: air oxidizing the alcohol reaction with air OR Due to volatility (of chemicals) IGNORE ...gas escaping ...HCl escaping		1

Question Number	Correct Answer	Reject	Mark
12 (a)(iv)	<p>First and second mark</p> <p>Phenolphthalein (1)</p> <p>From colourless to (pale) pink/red (1)</p> <p>ALLOW Other indicators with pK_{in} in range 7.5 – 10</p> <p>Some examples are:</p> <p>Thymol blue ((base)) (yellow to blue)</p> <p>Phenol red (yellow to red)</p> <p>Thymolphthalein (colourless to blue)</p> <p>Second mark depends on correct indicator except bromothymol blue, which is incorrect but very close to range so allow colour yellow to blue.</p> <p>Third mark Sodium ethanoate is (slightly) alkaline</p> <p>OR Ethanoic acid is a weak acid</p> <p>OR Phenolphthalein pH range coincides with vertical section of the pH/titration curve</p> <p>OR Titration of weak acid with strong base</p> <p>OR Neutralisation/equivalence point is at 8-10/ any number between 8 and 10.</p> <p>OR $pK_{in} \pm 1$ lies within vertical region (1)</p> <p>Third mark is independent</p>	<p>Litmus/universal indicator</p> <p>Pink to colourless</p> <p>Thymol blue (acid)</p> <p>Phenyl red Methyl red</p>	3

Question Number	Correct Answer	Reject	Mark
12 (b) (i)	$\text{CH}_3\text{COOH} + \text{CH}_3\text{CH}_2\text{OH} \rightleftharpoons \text{CH}_3\text{COOCH}_2\text{CH}_3 + \text{H}_2\text{O}$ <p>ALLOW</p> <p>Single arrow</p> <p>-CO₂H</p> <p>-C₂H₅</p> <p>Displayed formulae</p> <p>IGNORE state symbols even if incorrect</p>		1

Question Number	Correct Answer	Reject	Mark
12 (b) (ii)	<p>Volume of alkali reacting with ethanoic acid = $77.1 - 11.7 = 65.4 \text{ cm}^3$ (1)</p> <p>Moles of ethanoic acid = $\frac{65.4 \times 0.200}{1000}$ $= 0.01308 / 1.308 \times 10^{-2} \text{ (mol)}$ (1)</p> <p>Correct answer no working (2)</p> <p>Ignore SF except 1</p> <p>Allow internal TE for use of</p> <p>Moles of ethanoic acid = $\frac{77.1 \times 0.200}{1000}$ $= 0.01542 / 1.542 \times 10^{-2} \text{ (mol)}$ max(1)</p>		2

Question Number	Correct Answer	Reject	Mark
12 (b) (iii)	<p>Number of moles of ethanol =</p> <p>$0.01308 / 1.308 \times 10^{-2} \text{ (mol)}$</p> <p>TE same as (ii)</p>		1

Question Number	Correct Answer	Reject	Mark
12 (b)(iv)	Number of moles of ethyl ethanoate $= 0.0400 - 0.01308 = 0.02692$ (mol) Allow TE from (ii)/(iii) for example 0.01542 gives 0.02458		1

Question Number	Correct Answer	Reject	Mark
12 (b)(v)	$K_c = \frac{[\text{CH}_3\text{CO}_2\text{CH}_2\text{CH}_3][\text{H}_2\text{O}]}{[\text{CH}_3\text{CO}_2\text{H}][\text{CH}_3\text{CH}_2\text{OH}]}$ <p style="text-align: right;">(1)</p> $= \frac{0.02692 \times 0.02692}{0.01308 \times 0.01308}$ $= 4.23579 = 4.24$ <p style="text-align: right;">(1)</p> Ignore SF except one Allow TE from (ii), (iii) and (iv) for example 0.01542 etc gives 2.54 No TE for incorrect expression of K_c		2

Question Number	Correct Answer	Reject	Mark
12 (b)(vi)	The units cancel OR There are the same numbers of moles of reactants and products		1

Question Number	Correct Answer	Reject	Mark
12 (b)(vii)	(Concentrated) hydrochloric acid contains water		1

Question Number	Correct Answer	Reject	Mark
12 (c) (i)	First test tube esterification OR addition/elimination ALLOW Condensation (1) Second test tube (acid) hydrolysis (1) Two fully correct answers in wrong order (1) max	Alkaline hydrolysis followed by acidification	2

Question Number	Correct Answer	Reject	Mark
12 (c) (ii)	The values are the same within experimental error OR The values are concordant ALLOW The values are similar (1) The equilibrium can be approached from either direction OR The reaction is reversible OR Any comment relating equilibrium to reversibility IGNORE Dynamic equilibrium OR Rate of reverse reaction = rate of forward reaction (1)	Just...the same	2

Question Number	Correct Answer	Reject	Mark
12 (c) (iii)	(Acid) catalyst (makes it faster) OR Provides H ⁺ (as a catalyst) OR Protonates... OR Protonating agent... OR Donates protons OR Increases H ⁺ concentration	Initiates Reacts with... Protates	1

Total for Section C = 21 marks

Total for Paper = 90 marks

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE