




1.

Figure 1



A particle of weight  $24\text{ N}$  is held in equilibrium by two light inextensible strings. One string is horizontal. The other string is inclined at an angle of  $30^\circ$  to the horizontal, as shown in Figure 1. The tension in the horizontal string is  $Q$  newtons and the tension in the other string is  $P$  newtons. Find

(a) the value of  $P$ , (3)

(b) the value of  $Q$ . (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









4. A particle  $P$  of mass  $0.3\text{ kg}$  is moving with speed  $u\text{ m s}^{-1}$  in a straight line on a smooth horizontal table. The particle  $P$  collides directly with a particle  $Q$  of mass  $0.6\text{ kg}$ , which is at rest on the table. Immediately after the particles collide,  $P$  has speed  $2\text{ m s}^{-1}$  and  $Q$  has speed  $5\text{ m s}^{-1}$ . The direction of motion of  $P$  is reversed by the collision. Find

(a) the value of  $u$ , (4)

(b) the magnitude of the impulse exerted by  $P$  on  $Q$ . (2)

Immediately after the collision, a constant force of magnitude  $R$  newtons is applied to  $Q$  in the direction directly opposite to the direction of motion of  $Q$ . As a result  $Q$  is brought to rest in  $1.5\text{ s}$ .

(c) Find the value of  $R$ . (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





5. A ball is projected vertically upwards with speed  $21\text{ m s}^{-1}$  from a point  $A$ , which is  $1.5\text{ m}$  above the ground. After projection, the ball moves freely under gravity until it reaches the ground. Modelling the ball as a particle, find

(a) the greatest height above  $A$  reached by the ball, (3)

(b) the speed of the ball as it reaches the ground, (3)

(c) the time between the instant when the ball is projected from  $A$  and the instant when the ball reaches the ground. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









Leave blank

**Question 6 continued**

Lined writing area for the answer to Question 6, consisting of multiple horizontal lines.

**(Total 14 marks)**

**Q6**

--	--



7.

Figure 4




Figure 4 shows two particles  $P$  and  $Q$ , of mass 3 kg and 2 kg respectively, connected by a light inextensible string. Initially  $P$  is held at rest on a fixed smooth plane inclined at  $30^\circ$  to the horizontal. The string passes over a small smooth light pulley  $A$  fixed at the top of the plane. The part of the string from  $P$  to  $A$  is parallel to a line of greatest slope of the plane. The particle  $Q$  hangs freely below  $A$ . The system is released from rest with the string taut.

(a) Write down an equation of motion for  $P$  and an equation of motion for  $Q$ . (4)

(b) Hence show that the acceleration of  $Q$  is  $0.98 \text{ m s}^{-2}$ . (2)

(c) Find the tension in the string. (2)

(d) State where in your calculations you have used the information that the string is inextensible. (1)

On release,  $Q$  is at a height of 0.8 m above the ground. When  $Q$  reaches the ground, it is brought to rest immediately by the impact with the ground and does not rebound. The initial distance of  $P$  from  $A$  is such that in the subsequent motion  $P$  does not reach  $A$ . Find

(e) the speed of  $Q$  as it reaches the ground, (2)

(f) the time between the instant when  $Q$  reaches the ground and the instant when the string becomes taut again. (5)

---



---



---



---



---



---



Leave  
blank

**Question 7 continued**

Lined area for writing the answer to Question 7.

**Q7**

**(Total 16 marks)**

**TOTAL FOR PAPER: 75 MARKS**

**END**

