







3. A particle  $P$  of mass  $0.4 \text{ kg}$  moves under the action of a single constant force  $\mathbf{F}$  newtons. The acceleration of  $P$  is  $(6\mathbf{i} + 8\mathbf{j}) \text{ m s}^{-2}$ . Find

(a) the angle between the acceleration and  $\mathbf{i}$ , (2)

(b) the magnitude of  $\mathbf{F}$ . (3)

At time  $t$  seconds the velocity of  $P$  is  $\mathbf{v} \text{ m s}^{-1}$ . Given that when  $t = 0$ ,  $\mathbf{v} = 9\mathbf{i} - 10\mathbf{j}$ ,

(c) find the velocity of  $P$  when  $t = 5$ . (3)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



4. A car is moving along a straight horizontal road. The speed of the car as it passes the point  $A$  is  $25\text{ m s}^{-1}$  and the car maintains this speed for  $30\text{ s}$ . The car then decelerates uniformly to a speed of  $10\text{ m s}^{-1}$ . The speed of  $10\text{ m s}^{-1}$  is then maintained until the car passes the point  $B$ . The time taken to travel from  $A$  to  $B$  is  $90\text{ s}$  and  $AB = 1410\text{ m}$ .

(a) Sketch, in the space below, a speed-time graph to show the motion of the car from  $A$  to  $B$ .

(2)

(b) Calculate the deceleration of the car as it decelerates from  $25\text{ m s}^{-1}$  to  $10\text{ m s}^{-1}$ .

(7)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



Leave  
blank

**Question 4 continued** \_\_\_\_\_


Lined area for writing the answer to Question 4 continued.

**(Total 9 marks)**

Q4



5.



**Figure 1**

Two forces **P** and **Q** act on a particle at a point *O*. The force **P** has magnitude 15 N and the force **Q** has magnitude *X* newtons. The angle between **P** and **Q** is  $150^\circ$ , as shown in Figure 1. The resultant of **P** and **Q** is **R**.

Given that the angle between **R** and **Q** is  $50^\circ$ , find

(a) the magnitude of **R**, **(4)**

(b) the value of *X*. **(5)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



6.




Figure 2

A plank  $AB$  has mass  $12\text{ kg}$  and length  $2.4\text{ m}$ . A load of mass  $8\text{ kg}$  is attached to the plank at the point  $C$ , where  $AC = 0.8\text{ m}$ . The loaded plank is held in equilibrium, with  $AB$  horizontal, by two vertical ropes, one attached at  $A$  and the other attached at  $B$ , as shown in Figure 2. The plank is modelled as a uniform rod, the load as a particle and the ropes as light inextensible strings.

- (a) Find the tension in the rope attached at  $B$ . (4)

The plank is now modelled as a non-uniform rod. With the new model, the tension in the rope attached at  $A$  is  $10\text{ N}$  greater than the tension in the rope attached at  $B$ .

- (b) Find the distance of the centre of mass of the plank from  $A$ . (6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


---







7.

**Figure 3**

A package of mass 4 kg lies on a rough plane inclined at  $30^\circ$  to the horizontal. The package is held in equilibrium by a force of magnitude 45 N acting at an angle of  $50^\circ$  to the plane, as shown in Figure 3. The force is acting in a vertical plane through a line of greatest slope of the plane. The package is in equilibrium on the point of moving up the plane. The package is modelled as a particle. Find

- (a) the magnitude of the normal reaction of the plane on the package, (5)
- (b) the coefficient of friction between the plane and the package. (6)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





8.




Figure 4

Two particles  $P$  and  $Q$ , of mass 2 kg and 3 kg respectively, are joined by a light inextensible string. Initially the particles are at rest on a rough horizontal plane with the string taut. A constant force  $\mathbf{F}$  of magnitude 30 N is applied to  $Q$  in the direction  $PQ$ , as shown in Figure 4. The force is applied for 3 s and during this time  $Q$  travels a distance of 6 m. The coefficient of friction between each particle and the plane is  $\mu$ . Find

- (a) the acceleration of  $Q$ , (2)
- (b) the value of  $\mu$ , (4)
- (c) the tension in the string. (4)
- (d) State how in your calculation you have used the information that the string is inextensible. (1)

When the particles have moved for 3 s, the force  $\mathbf{F}$  is removed.

- (e) Find the time between the instant that the force is removed and the instant that  $Q$  comes to rest. (4)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



