

Centre No.						Paper Reference					Surname	Initial(s)	
Candidate No.					<b>6</b>	<b>6</b>	<b>7</b>	<b>8</b>	<b>/</b>	<b>0</b>	<b>1</b>	Signature	

Paper Reference(s)

**6678/01**

# Edexcel GCE

## Mechanics M2

### Advanced/Advanced Subsidiary

Friday 28 January 2011 – Morning

Time: 1 hour 30 minutes

Examiner's use only

--	--	--

Team Leader's use only

--	--	--

Question Number	Leave Blank
1	
2	
3	
4	
5	
6	
7	
8	
Total	

**Materials required for examination**

Mathematical Formulae (Pink)

**Items included with question papers**

Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

**Instructions to Candidates**

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of *g* is required, take  $g = 9.8 \text{ m s}^{-2}$ .

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

**Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

**Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled.

You should show sufficient working to make your methods clear to the examiner.

Answers without working may not gain full credit.



*Turn over*



Leave  
blank

**Question 1 continued**

Handwriting practice lines consisting of 30 horizontal lines spaced evenly down the page.

**(Total 6 marks)**

**Q1**



Leave  
blank

2. A particle of mass 2 kg is moving with velocity  $(5\mathbf{i} + \mathbf{j}) \text{ m s}^{-1}$  when it receives an impulse of  $(-6\mathbf{i} + 8\mathbf{j}) \text{ N s}$ . Find the kinetic energy of the particle immediately after receiving the impulse.

**(5)**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---





3. A particle moves along the  $x$ -axis. At time  $t = 0$  the particle passes through the origin with speed  $8 \text{ m s}^{-1}$  in the positive  $x$ -direction. The acceleration of the particle at time  $t$  seconds,  $t \geq 0$ , is  $(4t^3 - 12t) \text{ m s}^{-2}$  in the positive  $x$ -direction.

Find

(a) the velocity of the particle at time  $t$  seconds, (3)

(b) the displacement of the particle from the origin at time  $t$  seconds, (2)

(c) the values of  $t$  at which the particle is instantaneously at rest. (3)

Ruled lines for student answers











4.




Figure 1

A box of mass 30 kg is held at rest at point  $A$  on a rough inclined plane. The plane is inclined at  $20^\circ$  to the horizontal. Point  $B$  is 50 m from  $A$  up a line of greatest slope of the plane, as shown in Figure 1. The box is dragged from  $A$  to  $B$  by a force acting parallel to  $AB$  and then held at rest at  $B$ . The coefficient of friction between the box and the plane is  $\frac{1}{4}$ . Friction is the only non-gravitational resistive force acting on the box. Modelling the box as a particle,

- (a) find the work done in dragging the box from  $A$  to  $B$ . (6)

The box is released from rest at the point  $B$  and slides down the slope. Using the work-energy principle, or otherwise,

- (b) find the speed of the box as it reaches  $A$ . (5)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









5.




Figure 2

The uniform L-shaped lamina  $ABCDEF$ , shown in Figure 2, has sides  $AB$  and  $FE$  parallel, and sides  $BC$  and  $ED$  parallel. The pairs of parallel sides are 9 cm apart. The points  $A$ ,  $F$ ,  $D$  and  $C$  lie on a straight line.

$AB = BC = 36$  cm,  $FE = ED = 18$  cm.  $\angle ABC = \angle FED = 90^\circ$ , and  $\angle BCD = \angle EDF = \angle EFD = \angle BAC = 45^\circ$ .

(a) Find the distance of the centre of mass of the lamina from

(i) side  $AB$ ,

(ii) side  $BC$ .

(7)

The lamina is freely suspended from  $A$  and hangs in equilibrium.

(b) Find, to the nearest degree, the size of the angle between  $AB$  and the vertical.

(3)

---



---



---



---



---



---



---



---



---



---









Leave blank

Question 5 continued

Lined area for writing answers to Question 5.

(Total 10 marks)

Q5

Turn over



6. [In this question, the unit vectors  $\mathbf{i}$  and  $\mathbf{j}$  are in a vertical plane,  $\mathbf{i}$  being horizontal and  $\mathbf{j}$  being vertically upwards.]




Figure 3

At time  $t = 0$ , a particle  $P$  is projected from the point  $A$  which has position vector  $10\mathbf{j}$  metres with respect to a fixed origin  $O$  at ground level. The ground is horizontal. The velocity of projection of  $P$  is  $(3\mathbf{i} + 5\mathbf{j}) \text{ m s}^{-1}$ , as shown in Figure 3. The particle moves freely under gravity and reaches the ground after  $T$  seconds.

- (a) For  $0 \leq t \leq T$ , show that, with respect to  $O$ , the position vector,  $\mathbf{r}$  metres, of  $P$  at time  $t$  seconds is given by

$$\mathbf{r} = 3t\mathbf{i} + (10 + 5t - 4.9t^2)\mathbf{j} \quad (3)$$

- (b) Find the value of  $T$ . (3)
- (c) Find the velocity of  $P$  at time  $t$  seconds ( $0 \leq t \leq T$ ). (2)

When  $P$  is at the point  $B$ , the direction of motion of  $P$  is  $45^\circ$  below the horizontal.

- (d) Find the time taken for  $P$  to move from  $A$  to  $B$ . (2)
- (e) Find the speed of  $P$  as it passes through  $B$ . (2)

---



---



---



---



---





Leave blank

Question 6 continued

---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---



---





7.




Figure 4

A uniform plank  $AB$ , of weight  $100\text{ N}$  and length  $4\text{ m}$ , rests in equilibrium with the end  $A$  on rough horizontal ground. The plank rests on a smooth cylindrical drum. The drum is fixed to the ground and cannot move. The point of contact between the plank and the drum is  $C$ , where  $AC = 3\text{ m}$ , as shown in Figure 4. The plank is resting in a vertical plane which is perpendicular to the axis of the drum, at an angle  $\alpha$  to the horizontal, where  $\sin \alpha = \frac{1}{3}$ . The coefficient of friction between the plank and the ground is  $\mu$ . Modelling the plank as a rod, find the least possible value of  $\mu$ .

(10)

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---









Leave  
blank

**Question 7 continued**

Handwriting lines for the answer to Question 7.

**Q7**

**(Total 10 marks)**







Leave  
blank

**Question 8 continued**

Lined area for writing the answer to Question 8 continued.

**(Total 13 marks)**

**Q8**

--	--

**TOTAL FOR PAPER: 75 MARKS**

**END**

