

1. A car of mass 1000 kg moves with constant speed $V \text{ m s}^{-1}$ up a straight road inclined at an angle θ to the horizontal, where $\sin \theta = \frac{1}{30}$. The engine of the car is working at a rate of 12 kW. The resistance to motion from non-gravitational forces has magnitude 500 N. Find the value of V .

(5)



Question 1 continued

Lined writing area for Question 1 continued, consisting of approximately 20 horizontal lines.

(Total 5 marks)

Q1



Leave blank

Question 2 continued

Lined area for writing the answer to Question 2.

(Total 8 marks)

Q2



Question 3 continued

Lined area for writing the answer to Question 3.

(Total 8 marks)

Q3



Leave blank

Question 4 continued

Lined area for writing the answer to Question 4.

Q4

(Total 7 marks)



5.




Figure 2

A particle P of mass 0.5 kg is projected from a point A up a line of greatest slope AB of a fixed plane. The plane is inclined at 30° to the horizontal and $AB = 2\text{ m}$ with B above A , as shown in Figure 2. The particle P passes through B with speed 5 m s^{-1} . The plane is smooth from A to B .

- (a) Find the speed of projection. **(4)**

The particle P comes to instantaneous rest at the point C on the plane, where C is above B and $BC = 1.5\text{ m}$. From B to C the plane is rough and the coefficient of friction between P and the plane is μ .

By using the work-energy principle,

- (b) find the value of μ . **(6)**



Leave
blank

Question 5 continued

Lined writing area for the answer to Question 5 continued.



Leave
blank

Question 5 continued

Lined area for writing the answer to Question 5. The page contains 25 horizontal lines for student response.

Q5

--	--

(Total 10 marks)



P 3 8 1 6 2 A 0 1 5 2 8

6. A particle P moves on the x -axis. The acceleration of P at time t seconds is $(t - 4) \text{ m s}^{-2}$ in the positive x -direction. The velocity of P at time t seconds is $v \text{ m s}^{-1}$. When $t = 0, v = 6$.

Find

(a) v in terms of t , (4)

(b) the values of t when P is instantaneously at rest, (3)

(c) the distance between the two points at which P is instantaneously at rest. (4)



Leave
blank

Question 6 continued

Blank lined area for writing the answer to Question 6.



Leave
blank

Question 6 continued

Handwriting area with multiple horizontal lines.

(Total 11 marks)

Q6

--	--



7.




Figure 3

A uniform rod AB , of mass $3m$ and length $4a$, is held in a horizontal position with the end A against a rough vertical wall. One end of a light inextensible string BD is attached to the rod at B and the other end of the string is attached to the wall at the point D vertically above A , where $AD = 3a$. A particle of mass $3m$ is attached to the rod at C , where $AC = x$. The rod is in equilibrium in a vertical plane perpendicular to the wall as shown in Figure 3. The tension in the string is $\frac{25}{4}mg$.

Show that

(a) $x = 3a$, (5)

(b) the horizontal component of the force exerted by the wall on the rod has magnitude $5mg$. (3)

The coefficient of friction between the wall and the rod is μ . Given that the rod is about to slip,

(c) find the value of μ . (5)



Question 7 continued

Lined writing area for Question 7 continued.



Leave
blank

Question 7 continued

Lined area for writing the answer to Question 7.

Q7

--	--

(Total 13 marks)



8. A particle is projected from a point *O* with speed *u* at an angle of elevation α above the horizontal and moves freely under gravity. When the particle has moved a horizontal distance *x*, its height above *O* is *y*.

(a) Show that

$$y = x \tan \alpha - \frac{gx^2}{2u^2 \cos^2 \alpha} \tag{4}$$

A girl throws a ball from a point *A* at the top of a cliff. The point *A* is 8 m above a horizontal beach. The ball is projected with speed 7 m s⁻¹ at an angle of elevation of 45°. By modelling the ball as a particle moving freely under gravity,

(b) find the horizontal distance of the ball from *A* when the ball is 1 m above the beach. (5)

A boy is standing on the beach at the point *B* vertically below *A*. He starts to run in a straight line with speed *v* m s⁻¹, leaving *B* 0.4 seconds after the ball is thrown.

He catches the ball when it is 1 m above the beach.

(c) Find the value of *v*. (4)

Horizontal lines for working space.



Leave
blank

Question 8 continued

Lined area for writing the answer to Question 8.



