

Leave blank

Question 1 continued

Area containing horizontal lines for writing the answer to Question 1.

(Total 5 marks)

Q1



2.




Figure 1

The uniform lamina $OABCD$, shown in Figure 1, is formed by removing the triangle OAD from the square $ABCD$ with centre O . The square has sides of length $2a$.

(a) Show that the centre of mass of $OABCD$ is $\frac{2}{9}a$ from O . (4)

The mass of the lamina is M . A particle of mass kM is attached to the lamina at D to form the system S . The system S is freely suspended from A and hangs in equilibrium with AO vertical.

(b) Find the value of k . (4)



Leave
blank

Question 2 continued

Lined writing area for the answer to Question 2.

(Total 8 marks)

Q2



3. A particle P of mass 0.75 kg is moving with velocity $4\mathbf{i}\text{ m s}^{-1}$ when it receives an impulse $(6\mathbf{i} + 6\mathbf{j})\text{ N s}$. The angle between the velocity of P before the impulse and the velocity of P after the impulse is θ° .

Find

(a) the value of θ , (5)

(b) the kinetic energy gained by P as a result of the impulse. (3)



Question 3 continued

[Lined area for writing answer]



Question 3 continued

Lined writing area consisting of multiple horizontal lines for student responses.



Leave blank

Question 3 continued

Lined writing area for the answer to Question 3.

(Total 8 marks)

Q3



4. A ladder AB , of weight W and length $2l$, has one end A resting on rough horizontal ground. The other end B rests against a rough vertical wall. The coefficient of friction between the ladder and the wall is $\frac{1}{3}$. The coefficient of friction between the ladder and the ground is μ . Friction is limiting at both A and B . The ladder is at an angle θ to the ground, where $\tan \theta = \frac{5}{3}$. The ladder is modelled as a uniform rod which lies in a vertical plane perpendicular to the wall.

Find the value of μ .

(9)



Leave blank

Question 4 continued

Handwriting practice area with 24 horizontal lines.



Question 4 continued


Handwriting lines for the answer to Question 4. The page contains 28 horizontal lines for writing.

Q4

(Total 9 marks)



7.

**Figure 3**

At time $t = 0$, a particle is projected from a fixed point O on horizontal ground with speed $u \text{ m s}^{-1}$ at an angle θ° to the horizontal. The particle moves freely under gravity and passes through the point A when $t = 4 \text{ s}$. As it passes through A , the particle is moving upwards at 20° to the horizontal with speed 15 m s^{-1} , as shown in Figure 3.

- (a) Find the value of u and the value of θ . (7)

At the point B on its path the particle is moving downwards at 20° to the horizontal with speed 15 m s^{-1} .

- (b) Find the time taken for the particle to move from A to B . (2)

The particle reaches the ground at the point C .

- (c) Find the distance OC . (3)



Leave blank

Question 7 continued

Lined writing area consisting of multiple horizontal lines for student response.

(Total 12 marks)

Q7

--	--



P 4 4 8 3 7 A 0 2 7 3 2

8. Three identical particles P , Q and R , each of mass m , lie in a straight line on a smooth horizontal plane with Q between P and R . Particles P and Q are projected directly towards each other with speeds $4u$ and $2u$ respectively, and at the same time particle R is projected along the line away from Q with speed $3u$. The coefficient of restitution between each pair of particles is e . After the collision between P and Q there is a collision between Q and R .

(a) Show that $e > \frac{2}{3}$ (7)

It is given that $e = \frac{3}{4}$

(b) Show that there will not be a further collision between P and Q . (6)



