

Please check the examination details below before entering your candidate information

Candidate surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Tuesday 21 January 2020

Afternoon (Time: 1 hour 30 minutes)

Paper Reference **WME01/01**

Mathematics

International Advanced Subsidiary/Advanced Level
Mechanics M1

You must have:

Mathematical Formulae and Statistical Tables (Blue), calculator

Total Marks

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear.
Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either 2 significant figures or 3 significant figures.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

P61293A

©2020 Pearson Education Ltd.

1/1/1



Pearson

Leave blank

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 3 continued

Lined writing area for the answer to Question 3.



Leave blank

Question 3 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Handwriting practice area consisting of 30 horizontal lines.

(Total 11 marks)

Q3

--	--



4.




Figure 2

Two identical small rings, A and B , each of mass m , are threaded onto a rough horizontal wire. The rings are connected by a light inextensible string. A particle C of mass $3m$ is attached to the midpoint of the string. The particle C hangs in equilibrium below the wire with angle $BAC = \beta$, as shown in Figure 2.

The tension in each of the parts, AC and BC , of the string is T

(a) By considering particle C , find T in terms of m , g and β (2)

(b) Find, in terms of m and g , the magnitude of the normal reaction between the wire and A . (3)

The coefficient of friction between each ring and the wire is $\frac{4}{5}$

The two rings, A and B , are on the point of sliding along the wire towards each other.

(c) Find the value of $\tan\beta$ (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 4 continued

DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA
DO NOT WRITE IN THIS AREA

Lined writing area for the answer to Question 4.



5. A car travels at a constant speed of 40 m s^{-1} in a straight line along a horizontal racetrack. At time $t = 0$, the car passes a motorcyclist who is at rest. The motorcyclist immediately sets off to catch up with the car.

The motorcyclist accelerates at 4 m s^{-2} for 15 s and then accelerates at 1 m s^{-2} for a further T seconds until he catches up with the car.

(a) Sketch, on the same axes, the speed-time graph for the motion of the car and the speed-time graph for the motion of the motorcyclist, from time $t = 0$ to the instant when the motorcyclist catches up with the car. (2)

At the instant when $t = t_1$ seconds, the car and the motorcyclist are moving at the same speed.

(b) Find the value of t_1 (2)

(c) Show that $T^2 + kT - 300 = 0$, where k is a constant to be found. (6)



Question 6 continued

Lined area for writing the answer to Question 6. The area contains 30 horizontal lines for student input.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Q6

(Total 11 marks)

--	--



P 6 1 2 9 3 A 0 1 9 2 4

7.




Figure 3

A particle A of mass 4 kg is held at rest on a rough horizontal table. Particle A is attached to one end of a string that passes over a pulley P . The pulley is fixed at the edge of the table. The other end of the string is attached to a particle B , of mass 3 kg , which hangs freely below P .

The part of the string from A to P is perpendicular to the edge of the table and A , P and B all lie in the same vertical plane.

The string is modelled as being light and inextensible and the pulley is modelled as being small, smooth and light.

The system is released from rest with the string taut. At the instant of release, A is 2 m from the edge of the table and B is 1.4 m above a horizontal floor, as shown in Figure 3.

After descending with constant acceleration for 2 seconds, B hits the floor and does not rebound.

- (a) Show that the acceleration of A before B hits the floor is 0.7 m s^{-2} (2)
- (b) State which of the modelling assumptions you have used in order to answer part (a). (1)
- (c) Find the magnitude of the resultant force exerted on the pulley by the string. (4)
- The coefficient of friction between A and the table is μ .
- (d) Find the value of μ . (6)
- (e) Determine, by calculation, whether or not A reaches the pulley. (5)



Leave blank

Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the answer to Question 7.



