Please check the examination details b	elow before entering your candidate information
Candidate surname	Other names
Pearson Edexcel International Advanced Level	entre Number Candidate Number
Wednesday 8 J	lanuary 2020
Morning (Time: 1 hour 30 minutes)	Paper Reference WMA11/01
Mathematics	
International Advanced S Pure Mathematics P1	Subsidiary/Advanced Level
You must have: Mathematical Formulae and Statisti	cal Tables (Lilac), calculator

Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use black ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 11 questions in this question paper. The total mark for this paper is 75.
- The marks for each question are shown in brackets
- use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

 Turn over







1. Find, in simplest form,	$\int \left(\frac{8x^3}{3} - \frac{1}{2\sqrt{x}} - 5\right) \mathrm{d}x$	
		(4)

Question 1 continued	blank
	Q1
(Total 4 marks)	



2. Given $y = 3^x$, express each of the following in terms of y. Write each expression in its simplest form.



(1)

Leave blank

(b)
$$\frac{1}{3^{x-2}}$$

(2)

(c)	81	
(0)	9^{2-3x}	

(2)





Question 2 continued	Leave blank
	Q2
(Total 5 marks)	



3.




Figure 1

Figure 1 shows part of the curve with equation $y = x^2 + 3x - 2$

The point P(3,16) lies on the curve.

(a) Find the gradient of the tangent to the curve at P.

(2)

The point Q with x coordinate 3 + h also lies on the curve.

(b) Find, in terms of h, the gradient of the line PQ. Write your answer in simplest form. (3)

(c) Explain briefly the relationship between the answer to (b) and the answer to (a). (1)

6



Question 3 continued	blank
(Total 6 marks)	Q3



4.




Figure 2

Figure 2 shows the plan view of a house ABCD and a lawn APCDA.

ABCD is a rectangle with $AB = 16 \,\mathrm{m}$.

APCOA is a sector of a circle centre O with radius 12 m.

The point O lies on the line DC, as shown in Figure 2.

(a) Show that the size of angle AOD is 1.231 radians to 3 decimal places.

(2)

The lawn APCDA is shown shaded in Figure 2.

(b) Find the area of the lawn, in m², to one decimal place.

(4)

(c) Find the perimeter of the lawn, in metres, to one decimal place.

(3)



	Leav blanl
Question 4 continued	biani
Question 4 continued	



	Leave
	blank
Question 4 continued	
	Q4
(Total 9 marks)	
·	



(a) Find, using algebra, all solutions of

$$20x^3 - 50x^2 - 30x = 0$$

(3)

Leave blank

(b) Hence find all real solutions of

$$20(y+3)^{\frac{3}{2}} - 50(y+3) - 30(y+3)^{\frac{1}{2}} = 0$$

(4)

Question 5 continued	blank
	Q5
(Total 7 marks)	



The line l_1 has equation 3x - 4y + 20 = 0The line l_2 cuts the x-axis at R(8,0) and is parallel to l_1 (a) Find the equation of l_2 , writing your answer in the form ax + by + c = 0, where a, band c are integers to be found. The line l_1 cuts the x-axis at P and the y-axis at Q. Given that PQRS is a parallelogram, find (b) the area of PQRS, **(3)** (c) the coordinates of S. **(2)**



	Leave
	blank
Overting Countries d	Oldin
Question 6 continued	
	1



uestion 6 continued	

	Leave blank
Question 6 continued	
	Q6
(Total 8 marks)	
(



7.




Figure 3 shows part of the curve C_1 with equation $y = 3\sin x$, where x is measured in degrees.

The point P and the point Q lie on C_1 and are shown in Figure 3.

- (a) State
 - (i) the coordinates of P,
 - (ii) the coordinates of Q.

(3)

A different curve C_2 has equation $y = 3\sin x + k$, where k is a constant.

The curve C_2 has a maximum y value of 10

The point R is the minimum point on C_2 with the smallest positive x coordinate.

(b) State the coordinates of R.

(2)

Question 7 continued	Leave
	Q7
(Total 5 marks)	



8.	The straight line <i>l</i> has equation $y = k(2x - 1)$, where <i>k</i> is a constant.	
	The curve C has equation $y = x^2 + 2x + 11$	
	Find the set of values of k for which l does not cross or touch C .	(6)

Question 8 continued	blank
	Q8
(Total 6 marks)	



In this question you must show all stages of your working. 9. Solutions relying on calculator technology are not acceptable. A curve has equation $y = \frac{4x^2 + 9}{2\sqrt{x}} \qquad x > 0$ Find the x coordinate of the point on the curve at which $\frac{dy}{dx} = 0$ **(6)**



	Leave
	blank
Question 9 continued	
	Q9
(Total 6 marks)	



10. The curve C_1 has equation y = f(x), where

$$f(x) = (4x - 3)(x - 5)^2$$

- (a) Sketch C_1 showing the coordinates of any point where the curve touches or crosses the coordinate axes.
 - **(3)**

- (b) Hence or otherwise
 - (i) find the values of x for which $f\left(\frac{1}{4}x\right) = 0$
 - (ii) find the value of the constant p such that the curve with equation y = f(x) + p passes through the origin.

(2)

A second curve C_2 has equation y = g(x), where g(x) = f(x + 1)

- (c) (i) Find, in simplest form, g(x). You may leave your answer in a factorised form.
 - (ii) Hence, or otherwise, find the y intercept of curve C_2

(3)





Question 10 continued	blank
	Q10
(Total 8 marks)	



11. A curve has equation y = f(x), where

$$f''(x) = \frac{6}{\sqrt{x^3}} + x \qquad x > 0$$

The point P(4, -50) lies on the curve.

Given that f'(x) = -4 at P,

(a) find the equation of the normal at P, writing your answer in the form y = mx + c, where m and c are constants,

(3)

(b) find f(x).

(8)

	Leave
	blank
Question 11 continued	

