



Mark Scheme (Results)

October 2018

Pearson Edexcel International Advanced Level
in Statistics S1 (WST01/01)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2018

Publications Code WST01_01_1810_MS

All the material in this publication is copyright

© Pearson Education Ltd 2018

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

EDEXCEL IAL MATHEMATICS


General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
 6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
 7. Ignore wrong working or incorrect statements following a correct answer.

Question Number	Scheme	Marks
1.(a)	$S_{tt} = 1828 - \frac{(136)^2}{12} = 286.6\dots$ o.e. awrt 287	M1A1 (2)
(b)	$r = \frac{S_{ht}}{\sqrt{S_{tt}S_{hh}}} = \frac{-236}{\sqrt{286.6\dots \times 297}}$ or $\frac{-236}{\sqrt{85140}}$; = -0.8088... awrt -0.809	M1A1 (2)
(c)	Temperature decreases as height increases.	B1ft (1)
(d)	$b = \frac{S_{ht}}{S_{hh}} = \frac{-236}{297}$ (= -0.7946...) $a = \bar{t} - b\bar{h} = 11.3\dots + 0.7946 \times 9.33\dots = 18.7497\dots$ $t = 18.7 - 0.795h$	M1 M1 A1 (3)
(e)	$t = 18.7 - 0.795 \times 5 = 14.7$	M1 A1 (2)
(f)	Unreliable as the data is from France not South Africa	B1 (1) Total 11
Notes		
Correct numerical answers in (a), (b), (d) or (e) score all the marks for that part.		
(a)	M1 Correct expression for S_{tt} A1 awrt 287 allow exact fractions e.g. $\frac{860}{3}$ or $286\frac{2}{3}$	
(b)	M1 for attempt at correct formula, values must be substituted. Allow $\frac{-236}{\sqrt{287 \times 297}}$ A1 awrt - 0.809 (allow - 0.808 from a correct expression with 287 used)	
(c)	B1ft for a comment in context. Must see “height” (or h) and “temperature” (or t) mentioned Allow “as the temperature increases the height above sea level decreases” (o.e.) NB If $ r > 1$ score B0 in (c) Saying “sea level increases” (o.e.) is B0	
(d)	M1 Correct expression for b . M1 Allow 11.3... - “their $b \times 9.33\dots$ ” [$a = \frac{16706}{891}$ scores M1 but A0] A1 $t = (18.75$ or awrt 18.7) - (awrt 0.795) h [No fractions and no x, y]	
(e)	M1 substitute $h = 5$ or 500 into <u>their</u> regression line A1 answer in range [14.7 , 14.8] (condone coming from y, x equation)	
(f)	B1 unreliable with a reason. [Use of 500 in (e) <u>and</u> stating “out of range” is B0] Must mention France or (S) Africa and at least imply the other	

Question Number	Scheme	Marks
<p>2.(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>$Q_2 = 54$ $Q_1 = 45$ $Q_3 = 59$</p> <p>Upper limit = $59 + 1.5 \times 14 = 80$ Lower limit = $45 - 1.5 \times 14 = 24$ Outlier 84</p>  <p>Any two from: The females are heavier than the males (on average). The males have lower median than females. The males have a smaller IQR than the females. The females have a greater range than males. [Comments just about skewness are B0]</p>	<p>B1 B1 B1 (3)</p> <p>M1 A1 A1ft (3)</p> <p>B1 B1 B1 (3)</p> <p>B1 B1 (2)</p>
Notes		Total 11
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>1st B1 for $Q_2 = 54$ 2nd B1 for $Q_1 = 45$ 3rd B1 for $Q_3 = 59$</p> <p>M1 correct expression for either limit ft their values in (a) 1st A1 80 and 24 2nd A1ft all outliers identified using their limits (must be stated in (b))</p> <p>1st B1 Box with whiskers drawn and Q_2 and quartiles ft from (a), condone 2 whiskers on RHS 2nd B1 For only one lower whisker to 32 and no outliers 3rd B1 For upper whisker to 80 or 77 and an outlier at 84 NB If there are whiskers at both 77 and 80 it is 3rd B0</p> <p>1st B1 a correct comparison on location e.g. median or comment implying “on average” 2nd B1 a second correct comparison on spread e.g. range or IQR (greater spread is B0)</p>	

Question Number	Scheme	Marks
<p>3(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>Width = 3 cm 1cm^2 represents 2 cars <u>or</u> 0.5cm^2 represents 1 car <u>or</u> their $h \times w = 6$ <u>or</u> area = 6 Height = $\frac{6}{3} = 2$ cm</p> <p>Median = $(2) + \frac{30-28}{15} \times 2$ <u>or</u> $(2) + \frac{30.5-28}{15} \times 2$ (o.e.) = 2.266... (or 2.33...)</p> <p>$[\bar{t}] = \frac{182}{60} = 3.03...$ $[\sigma_t] = \sqrt{\frac{883}{60} - \bar{t}^2} = \sqrt{5.5155...}$ = 2.3485... ($s = 2.3683...$)</p> <p>Mean > median Positive skew</p> <p>[75 mins = 1.25 hours] $(> 75 \text{ mins}) = 5 + 12 + 15 + \frac{3}{4} \times 18 = 45.5$ <u>or</u> $(< 75) = 10 + \frac{1}{4} \times 18$ <u>or</u> $28 - \frac{3}{4} \times 18$ $P(T > 1.25) = \frac{45.5}{60}$ <u>or</u> e.g. $1 - \frac{14.5}{60}$ 0.7583... awrt 0.758</p>	<p>B1 M1 A1 (3)</p> <p>M1 A1 (2)</p> <p>B1</p> <p>M1 A1 (3)</p> <p>B1ft dB1 (2)</p> <p>M1 M1 A1 (3)</p>
Notes		Total 13
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>NB</p>	<p>B1 3 only M1 may be implied by correct height A1 correct height of 2(cm) and correct width of 3 (cm)</p> <p>M1 for any correct equation leading to correct fraction as part of $m = \dots$ or $(m - [2]) = \dots$ Ignore incorrect end point and watch out for “working down” A1 awrt 2.27 allow exact fraction e.g. $\frac{34}{15}$ (allow awrt 2.33 [or $\frac{7}{3}$] if $n + 1$ used)</p> <p>B1 awrt 3.03 (allow exact fraction e.g. $\frac{91}{30}$) M1 A correct expression A1 awrt 2.35 or 2.37</p> <p>1st B1 ft their mean and median (Allow “larger frequencies at the start of table”) Do not allow comparison of quartiles unless correct values are seen (2sf comparisons) $Q_1 = 1.28$ or $\frac{23}{18}$ [$(n+1) = 1.29$] $Q_3 = 4.33$ or $\frac{13}{3}$ [$(n+1) = 4.42$] e.g. $2.1 > 0.99$ or $2.1 > 1.0$ 2nd dB1 dependent on previous B1 being awarded.</p> <p>1st M1 for a correct expression for no. of cars longer than 75 mins or shorter than 75 mins 2nd M1 $\frac{k}{60}$ where $44 \leq k < 46$ A1 awrt 0.758 allow $\frac{91}{120}$ (o.e.)</p> <p>Any use of the normal distribution is M0M0A0</p>	

Question Number	Scheme	Marks
<p>4.(a)</p> <p>(b)</p> <p>Ans only</p> <p>(c)</p> <p>(d)</p>	<p>0.13</p> <p>$P(A) \times P(C) = P(A \cap C)$</p> <p>$0.2 \times (0.08 + p) = 0.05$ or $P(C) = \frac{0.05}{0.10 + 0.05 + 0.01 + 0.04}$ or $\frac{0.05}{0.2}$ or 0.25</p> <p>$p = 0.17$</p> <p>$P(\text{no faults}) = 1 - (0.1 + 0.05 + 0.01 + 0.04 + 0.08 + 0.03 + "0.17")$</p> <p>or $1 - ["P(C)" + 0.10 + 0.05 + 0.08]$</p> <p>$q = \underline{\underline{0.52}}$</p> <p>They can get q without finding p so a correct answer to q scores 4/4</p> <p>$P(\text{Fault } B \text{ but not fault } C \mid \text{Has fault } A) = \frac{0.05}{0.2}$</p> <p>$= 0.25$</p> <p>$P(\text{exactly 2 defects}) = 0.12$ or $\frac{3}{25}$</p> <p>$P(\text{both have 2 defects}) = 0.12^2$</p> <p>$= \underline{\underline{0.0144}}$ or $\frac{9}{625}$</p>	<p>B1</p> <p>(1)</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>(4)</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>Total 10</p>
Notes		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>B1 0.13 only</p> <p>1st M1 using $P(A) \times P(C) = P(A \cap C)$ allow one addition error in $P(A)$ e.g. $P(A) = 0.11$</p> <p>1st A1 0.17 only</p> <p>2nd M1 $1 - (0.10 + 0.05 + 0.01 + 0.04 + 0.08 + 0.03 + "their 0.17")$ allow letter p for 0.17</p> <p>or $1 - ["P(C)" + 0.10 + 0.05 + 0.08]$ but need a value for $P(C)$ [M0A0M1A0 possible]</p> <p>2nd A1 0.52 only (correct answer of 0.52 with no incorrect working is 4/4)</p> <p>M1 for attempt at $P(B \cap C' \mid A)$ allow for $\frac{0.06}{0.2}$ or $\frac{0.05}{0.2}$ allow ft of their $P(A)$ used in part(b)</p> <p>A1 0.25</p> <p>B1 sight of 0.12 or $(0.05 + 0.03 + 0.04)$ only NB e.g. 0.12×2 is B1M0A0</p> <p>M1 $("0.12")^2$ where $0.1 < "0.12" < 0.2$</p> <p>May see attempt at $(0.05 + 0.03 + 0.04)^2$ multiplied out but must have ≥ 4 correct products</p> <p>A1 0.0144 (o.e.) (correct answer only scores 3/3)</p>	

Question Number	Scheme	Marks																		
<p>5. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	$k = \frac{2}{35}$ <table border="1" data-bbox="268 259 1294 421"> <tr> <td>x</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> </tr> <tr> <td>$P(X=x)$</td> <td>$\frac{3}{35}$</td> <td>$\frac{5}{35}$ or $\frac{1}{7}$</td> <td>$\frac{7}{35}$ or $\frac{1}{5}$</td> <td>$\frac{9}{35}$</td> <td>$\frac{11}{35}$</td> </tr> <tr> <td>$P(X=x)$</td> <td>0.08571..</td> <td>0.14285..</td> <td>0.2</td> <td>0.25714..</td> <td>0.31428..</td> </tr> </table> $\frac{5}{35} + \frac{7}{35} = \frac{12}{35}$ $E(X) = 1 \times \frac{3}{35} + 2 \times \frac{5}{35} + 3 \times \frac{7}{35} + 4 \times \frac{9}{35} + 5 \times \frac{11}{35} = \left[\frac{25}{7} \right]$ $E(X^2) = 1 \times \frac{3}{35} + 4 \times \frac{5}{35} + 9 \times \frac{7}{35} + 16 \times \frac{9}{35} + 25 \times \frac{11}{35} = \left[\frac{101}{7} \right]$ $\text{Var}(X) = \frac{101}{7} - \left(\frac{25}{7} \right)^2 ; = \frac{82}{49} \quad (\text{allow } 1.67 \sim 1.674)$ $\text{Var}(12 - 7X) = 7^2 \times \frac{82}{49} ; = \underline{\underline{82}}$ <p>$4X \leq Y$ when $X = 1, 4$ or 5, so probability = "$\frac{3}{35}$" + "$\frac{9}{35}$" + "$\frac{11}{35}$"</p> $= \frac{23}{35}$	x	1	2	3	4	5	$P(X=x)$	$\frac{3}{35}$	$\frac{5}{35}$ or $\frac{1}{7}$	$\frac{7}{35}$ or $\frac{1}{5}$	$\frac{9}{35}$	$\frac{11}{35}$	$P(X=x)$	0.08571..	0.14285..	0.2	0.25714..	0.31428..	<p>B1</p> <p>M1 A1 (3)</p> <p>M1 A1ft (2)</p> <p>M1</p> <p>M1</p> <p>M1; A1</p> <p>M1; A1 (6)</p> <p>M1; A1ft</p> <p>A1 (3)</p>
x	1	2	3	4	5															
$P(X=x)$	$\frac{3}{35}$	$\frac{5}{35}$ or $\frac{1}{7}$	$\frac{7}{35}$ or $\frac{1}{5}$	$\frac{9}{35}$	$\frac{11}{35}$															
$P(X=x)$	0.08571..	0.14285..	0.2	0.25714..	0.31428..															
Notes		Total 14																		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>Dist of Y</p> <p>F(x) = f(x)</p> <p>All in k</p> <p>Reverse Y</p>	<p>B1 may be implied by a correct probability</p> <p>M1 need x values each with a prob and at least one correct prob. (Allow probs in terms of k)</p> <p>A1 all values correct – accept decimals 3sf or better</p> <p>M1 “their $P(X=2)$” + “their $P(X=3)$”</p> <p>A1ft ft providing <1 Allow answer in $[0.3428, 0.343]$ or $6k$</p> <p>1st M1 using $\sum xP(X=x)$ or $\frac{25}{7}$ or $\frac{125}{2}k$ or $\sum yP(Y=y)$ or -13 (≥ 4 correct terms or ft)</p> <p>2nd M1 using $\sum x^2P(X=x)$ or $\frac{101}{7}$ or $\frac{505}{2}k$ or $\sum y^2P(Y=y)$ (≥ 4 correct terms or ft)</p> <p>3rd M1 using $\text{Var}(X) = E(X^2) - [E(X)]^2$ or $\text{Var}(Y) = E(Y^2) - [E(Y)]^2$</p> <p>1st A1 for a correct answer (allow 3sf) or for $E(Y^2) = 251$</p> <p>4th M1 $49 \times \text{Var}(X)$ or correct distribution for Y (ft probs from X)</p> <p>2nd A1 for 82 only</p> <p>M1 for $X = 1, 4$ or 5 [or $Y = 5, -16, -23$] and at least one correct ft probability.</p> <p>A1ft their "$\frac{3}{35}$"; + their "$\frac{9}{35}$" + their "$\frac{11}{35}$" providing sum is <1 (allow in terms of k)</p> <p>A1 cao (allow $\frac{23}{2}k$)</p> <table border="1" data-bbox="268 1805 1294 1928"> <tr> <td>y^2</td> <td>25</td> <td>4</td> <td>81</td> <td>256</td> <td>529</td> </tr> <tr> <td>y</td> <td>5</td> <td>-2</td> <td>-9</td> <td>-16</td> <td>-23</td> </tr> <tr> <td>$P(X=x)$</td> <td>$\frac{3}{35}$</td> <td>$\frac{5}{35}$ or $\frac{1}{7}$</td> <td>$\frac{7}{35}$ or $\frac{1}{5}$</td> <td>$\frac{9}{35}$</td> <td>$\frac{11}{35}$</td> </tr> </table> <p>Get $k = \frac{2}{85}$ Can award: (a) 0/3 (b) M1A1ft (c) M4A0 (d) M1A1ftA0</p> <p>Can award: (a) B0M1A0 (b) 2/2 in (c) M4A0 (d) M1A1ftA1</p> <p>May see $Y = 12 - 7(6 - X)$ used: in (c) can score M3 A0 probably zero in (d)</p>	y^2	25	4	81	256	529	y	5	-2	-9	-16	-23	$P(X=x)$	$\frac{3}{35}$	$\frac{5}{35}$ or $\frac{1}{7}$	$\frac{7}{35}$ or $\frac{1}{5}$	$\frac{9}{35}$	$\frac{11}{35}$	
y^2	25	4	81	256	529															
y	5	-2	-9	-16	-23															
$P(X=x)$	$\frac{3}{35}$	$\frac{5}{35}$ or $\frac{1}{7}$	$\frac{7}{35}$ or $\frac{1}{5}$	$\frac{9}{35}$	$\frac{11}{35}$															

Question Number	Scheme	Marks		
6. (a)	$P(L > 4.3) = P\left(Z > \frac{4.3 - 4.1}{0.125}\right)$ $= P(Z > 1.6) \text{ or } 1 - P(Z < 1.6) \text{ or } 1 - 0.9452$ $= 0.0548$	M1 M1 A1 (3)		
(b)	$P(3.9 < L < 4.3) = P(Z < 1.6) - P(Z < -1.6) \text{ or } 2(P(Z < 1.6) - 0.5)$ $= 0.9452 - 0.0548 \qquad = 2(0.9452 - 0.5)$ $= 0.8904 \qquad = 0.8904$	B1cso (1)		
(c)	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; vertical-align: top;"> Number of unusable bolts $= (1 - 0.89) \times 500 [= 55]$ Value of bolts = "445" × 9 + "55" × 1 profit = "445" × 9 + "55" × 1 - 500 × 5 Profit from bolts = 1560 pence </td> <td style="width: 50%; vertical-align: top;"> <u>Alternative</u> E(value of a bolt) = 0.89 × 9 + 0.11 × 1 E(profit per bolt) = 0.89 × 9 + 0.11 × 1 - 5 Profit = "3.12" × 500 Profit from bolts = 1560 pence </td> </tr> </table>	Number of unusable bolts $= (1 - 0.89) \times 500 [= 55]$ Value of bolts = "445" × 9 + "55" × 1 profit = "445" × 9 + "55" × 1 - 500 × 5 Profit from bolts = 1560 pence	<u>Alternative</u> E(value of a bolt) = 0.89 × 9 + 0.11 × 1 E(profit per bolt) = 0.89 × 9 + 0.11 × 1 - 5 Profit = "3.12" × 500 Profit from bolts = 1560 pence	M1oe M1oe M1oe A1 (4)
Number of unusable bolts $= (1 - 0.89) \times 500 [= 55]$ Value of bolts = "445" × 9 + "55" × 1 profit = "445" × 9 + "55" × 1 - 500 × 5 Profit from bolts = 1560 pence	<u>Alternative</u> E(value of a bolt) = 0.89 × 9 + 0.11 × 1 E(profit per bolt) = 0.89 × 9 + 0.11 × 1 - 5 Profit = "3.12" × 500 Profit from bolts = 1560 pence			
(d)	$\frac{4.198 - \mu}{\sigma} = 1.96 \quad \text{or} \quad 4.198 - \mu = 1.96\sigma \quad \text{oe}$ $\frac{4.065 - \mu}{\sigma} = -0.7 \quad \text{or} \quad 4.065 - \mu = -0.7\sigma \quad \text{oe}$ $0.133 = 2.66\sigma$ $\sigma = 0.05 \quad (\text{or awrt } 0.0500)$ $\mu = 4.1 \quad (\text{or awrt } 4.10)$	M1A1 A1 M1 A1 (6)		
(e)	The mean the same but the st. dev. decreased or $P(3.9 < L < 4.3)$ increased. So the profit will increase NB Use of + 0.7 in (c) → $\mu = 3.99, \sigma = 0.106$, prob ≈ 0.80 → profit down	B1ft dB1ft (2) Total 16		
Notes				
(a)	1 st M1 standardising. Allow use of 0.125^2 2 nd M1 $1 - p$ $p > 0.8$ A1 awrt 0.0548			
(b)	B1cso sight of 0.8904 or better (calc: 0.8904014212...) or a correct subtraction			
(c)	1 st M1 $(1 - "0.89") \times 500$ <u>or</u> $0.89 \times 9 + 0.11 \times 1$ 2 nd M1 "445" × 9 + "55" <u>or</u> $0.89 \times 9 + 0.11 \times 1 - 5$ 3 rd M1 method for the profit <u>or</u> their "3.12" × 500 A1 for awrt £15.60 or 1560 pence(p) [need units]	SC think 55 scrap loses 1p 1 st M1 for sight of 55 B1 for answer of awrt 1450 p Score as: M1M0M0A1		
(d)	1 st M1 Forming either equation – must have z value but allow $\pm z$ where $ z > 0.6$ 1 st A1 correct equation $4.198 - \mu = 1.96\sigma$ - any form (or allow $z =$ awrt 1.960) 2 nd A1 correct equation $4.065 - \mu = -0.7\sigma$ - any form (or allow $z =$ awrt -0.700) 2 nd M1 eliminating μ or σ (method <u>seen</u> leading to equation in 1 variable) 3 rd A1 0.05 (or awrt 0.0500) 4th A1 4.1 (or awrt 4.10 dep on 1st or 2nd A1) NB Candidate who assumes $\mu = 4.1$ can get M1 A0 A0M1A0A1			
(e)	1 st B1ft if $\mu = 4.1$ then ft σ ; if $\mu < 3.9$ (allow any σ) <u>otherwise</u> need to see $P(3.9 < L < 4.3)$ calc If they have $\mu = 4.1$ in part (d) then don't need to state "mean the same" in part (e) 2 nd dB1ft therefore profit will increase (o.e.) [$\sigma < 0$ is B0B0]			

