



Mark Scheme (Results)

January 2021

Pearson Edexcel International GCSE
Mathematics A (4MA1)
Paper 1HR

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2021

Publications Code 4MA1_1HR_2101_MS

All the material in this publication is copyright

© Pearson Education Ltd 2021

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme.

Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.

- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

- **Types of mark**

- M marks: method marks
- A marks: accuracy marks
- B marks: unconditional accuracy marks (independent of M marks)

- **Abbreviations**

- cao – correct answer only
- ft – follow through
- isw – ignore subsequent working
- SC - special case
- oe – or equivalent (and appropriate)

- dep – dependent
- indep – independent
- awrt – answer which rounds to
- eoo – each error or omission

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.

- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the “correct” answer has been obtained from incorrect working, award 0 marks.

If a candidate misreads a number from the question. Eg. Uses 252 instead of 255; method marks may be awarded provided the question has not been simplified. Examiners should send any instance of a suspected misread to review. If there is a choice of methods shown, mark the method that leads to the answer on the answer line; where no answer is given on the answer line, award the lowest mark from the methods shown.

If there is no answer on the answer line then check the working for an obvious answer.

- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.

- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded to another.

International GCSE Maths				
Apart from questions 1, 8, 10, 11d, 12c, 14, 15ab, 17 (where the mark scheme states otherwise) the correct answer, unless clearly obtained by an incorrect method, should be taken to imply a correct method				
Question	Working	Answer	Mark	Notes
1	e.g. $\frac{16}{5}$ and $\frac{11}{6}$ or $\frac{96}{30}$ and $\frac{55}{30}$		3	M1 for two correct improper fractions
	e.g. $\frac{16^8}{5} \times \frac{11}{6^3}$ or $\frac{176}{30}$ or $\frac{5280}{900}$ oe			M1 correct cancelling or multiplication of numerators and denominators without cancelling
	e.g. $\frac{16}{5} \times \frac{11}{6} = \frac{176}{30} = \frac{88}{15} = 5\frac{13}{15}$ or $\frac{16}{5} \times \frac{11}{6} = \frac{176}{30} = 5\frac{26}{30} = 5\frac{13}{15}$ or $\frac{16^8}{5} \times \frac{11}{6^3} = \frac{88}{15} = 5\frac{13}{15}$ or $\frac{96}{30} \times \frac{55}{30} = \frac{5280}{900} = \frac{88}{15} = 5\frac{13}{15}$ NB: a student can show initially that $5\frac{13}{15} = \frac{88}{15}$ and they need to show that LHS = $\frac{88}{15}$	shown		A1 Dep on M2 for conclusion to $5\frac{13}{15}$ from correct working – either sight of the result of the multiplication e.g. $\frac{176}{30}$ must be seen and equated to $\frac{88}{15}$ or $5\frac{26}{30}$ or correct cancelling prior to the multiplication to $\frac{88}{15}$ NB: use of decimals scores no marks
				Total 3 marks

2	$a = 7$		4	B1
	$\frac{b + \text{their } a}{2} = 8.5$ oe or $b = 10$			M1 ft their value of a or for setting up an equation for b or $b = 10$
	$\frac{\text{their } a + \text{their } a + \text{their } b + c}{4} = 9$ oe or $(c =) 9 \times 4 - (2 \times \text{their } a + \text{their } b)$ oe			M1 for a calculation involving c using their values or for a calculation leading to c using their values
		7, 10, 12		A1
				Total 4 marks

3	a	Correct number line	2	B2 for a fully correct number line e.g. shaded circle at -2 , unshaded circle at 1 and a line drawn between them B1 for a shaded circle at -2 or an unshaded circle at 1 or circles at -2 and 1 with line in between but shading incorrect
	b	$-3, -2, -1, 0, 1, 2$	2	B2 fully correct values with no extras B1 for 5 correct values and none incorrect or all 6 correct values with no more than one additional incorrect value
				Total 4 marks

4	3.4 or $\frac{17}{5}$ or $3\frac{2}{5}$ or $3\frac{24}{60}$ or 204 oe		3	B1
	$433.5 \div 3.4$ or $433.5 \div \frac{17}{5}$ or $433.5 \div 3\frac{2}{5}$ or $\frac{433.5}{'204'} \times 60$ oe			M1 for use of speed = distance \div time Allow $433.5 \div 3.24$ (= 133.796...) for this mark only
		127.5		A1 oe allow 128
				Total 3 marks

5	a	$(x =) 270 \div (12 \times 5)$ (= 4.5) oe		3	M1
		$\pi \times '4.5'^2 \times 2 \times '4.5'$ (= 182.25π oe)			M1 ft dep on M1
			573		A1 accept 572 – 573
	b		1 000 000	1	B1 or $(1 \times) 10^6$ or (one or 1) million oe
					Total 4 marks

6	a	e.g. $A + 5z = \frac{c}{y}$ oe or $Ay = c - 5yz$ oe		2	M1 for a correct first step e.g. add $5z$ to both sides or multiply all terms by y
			$c = y(A + 5z)$		A1 oe
	b		1	1	B1
	c	$(x \pm 3)(x \pm 8)$		2	M1 or for $(x \pm a)(x \pm b)$ where $ab = 24$ or $a + b = -11$
			$(x - 3)(x - 8)$		A1
					Total 5 marks

7	$0.024 \times 50\,000 (= 1200)$ oe or $1.024 \times 50\,000 (= 51\,200)$ oe or $1.024^2 \times 50\,000 (= 52\,428.8)$ oe or $0.024 \times 50\,000 \times 3 (= 3600)$ oe $0.024 \times 50\,000 \times 3 + 50\,000 (= 53\,600)$ oe		3	M1	M2 for $50\,000 \times 1.024^3$
	$0.024 \times (50\,000 + '1200')$ (= 1228.8) oe and $0.024 \times (50\,000 + '1200' + '1228.8')$ (= 1258.2912) or '1200' + '1228.8' + '1258.2912' (= 3687.(0912)) or $1.024 \times '52\,428.8'$			M1 for completing method to find total amount in the account	
		53 687		A1 accept 53 687 – 53 688	
				accept $(1 + 0.024)$ or $\left(1 + \frac{2.4}{100}\right)$ as equivalent to 1.024 throughout	
					Total 3 marks

8	$(5 - 2) \times 180 \div 5 (= 108)$ or $360 \div 5 (= 72)$		5	M1	for method to find an interior or exterior angle of a pentagon
	$(6 - 2) \times 180 \div 6 (= 120)$ or $360 \div 6 (= 60)$			M1	for method to find an interior or exterior angle of a hexagon
	$360 - 108 - 120 (= 132)$ or $60 + 72 (= 132)$ or $(180 - '120') + (180 - '108')$			M1	dep on M2 for a correct method to find angle <i>EDI</i> using correct figures
	$360 - '72' - '60' - '132' (= 96)$			M1	for a complete method to find angle <i>x</i>
		96		A1	dep on correct working
				Note:	Angles may be seen on diagram throughout
Total 5 marks					

9	a		$2^6 \times 3 \times 11^4$	2	B2	oe, accept 2 811 072
					B1	for $2^a \times 3^b \times 11^c$ oe where two of <i>a</i> , <i>b</i> and <i>c</i> are correct
	b		$2^9 \times 3^5 \times 11^8$	2	B2	cao
					B1	for $2^a \times 3^b \times 11^c$ oe where two of <i>a</i> , <i>b</i> and <i>c</i> are correct or $2.666... \times 10^{13}$ or an equivalent expression for e.g. $2^2 \times 2^7 \times 3^5 \times 11^3 \times 11^5$
Total 4 marks						

10	$7^2 - (10 \div 2)^2 (= 24) \text{ or } \frac{\sin\left(\frac{1}{2}x\right)}{5} = \frac{\sin 90}{7} \text{ oe or}$ $\cos x = \frac{7^2 + 7^2 - 10^2}{2 \times 7 \times 7} \text{ oe or } \sin\left(\frac{1}{2}x\right) = \frac{5}{7} \text{ oe or } \cos y = \frac{5}{7} \text{ oe}$	5	M1 or use of sine rule or cosine rule to find angle (x) of the apex or angle y $\left(= 90 - \frac{1}{2}x\right)$
	$\sqrt{7^2 - (10 \div 2)^2} (= \sqrt{24} = 2\sqrt{6} = 4.898...) \text{ or}$ $(x =) 2 \times \sin^{-1}\left(\frac{5 \times \sin 90}{7}\right) (= 91.169...) \text{ oe or}$ $(x =) 2 \times \sin^{-1}\left(\frac{5}{7}\right) (= 91.169...) \text{ oe or}$ $(x =) \cos^{-1}\left(\frac{7^2 + 7^2 - 10^2}{2 \times 7 \times 7}\right) (= 91.169...) \text{ oe or}$ $(x =) 2\left(90 - \cos^{-1}\left(\frac{5}{7}\right)\right) (= 2(90 - 44.415)... = 91.169...)$ <p>Allow 5 from correct working</p>		M1 for complete method to find height of triangle or the angle (x) of the apex $\cos^{-1}\left(\frac{5}{7}\right) (= 44.415...) \text{ and}$ $5 \times \tan'44.415...' (= 4.898...) \text{ or}$ $7 \times \sin'44.415...' (= 4.898...)$ or $\sin^{-1}\left(\frac{5}{7}\right) (= 45.584...) \text{ and}$ $\frac{5}{\tan'45.584...' (= 4.898...) \text{ or}$ $7 \times \cos'45.584...' (= 4.898...)$
	<p>E.g.</p> $6 \times 10 + \frac{(10 \div 2) \times \sqrt{24}}{2} \times 2 (= 60 + 10\sqrt{6} = 84.494...) \text{ or}$ $5 \times (6 + 6 + \sqrt{24}) (= 60 + 10\sqrt{6} = 84.494...) \text{ or}$ $\left(\frac{1}{2} \times 7 \times 7 \times \sin'91.169...' + 10 \times 6\right) (= 60 + 10\sqrt{6} = 84.494...)$		M1 for method to find the total area of the pentagon allow answers in the range 84.49 – 85
	<p>E.g.</p> $'84.494' \div 16 (= 5.28...) \text{ or } (60 + 10\sqrt{6}) \div 16 (= 5.28...)$		M1 for method to find the number of tins required using their area
		6	A1 dep on at least M2
			Total 5 marks

11	a		8, 23, 40, 68, 101, 120	1	B1
	b			2	M1 ft from table for at least 5 points plotted correctly at end of interval or ft from sensible table for all 6 points plotted consistently within each interval in the freq table at the correct height
			Correct cf graph		A1 accept curve or line segments accept graph that is not joined to (0,0)
	c		17 – 20	1	B1 ft their cf graph
	d	E.g. Reading at 23 minutes (= <i>a</i>) and then $(120 - a) \div 120 \times 100$		2	M1 ft from their cf graph reading off at 23 minutes and a method to work out 120 minus this value as a percentage of 120
			25(%) – 29(%)		A1 ft from their cf graph dep on M1 seen
					Total 6 marks

12	a		$4e^5 f^3$	2	B2 (B1 for 2 out of 3 terms correct in a 3 term product)
	b	E.g. $\frac{3(2x+1)+4(x-2)}{12}$ or $\frac{3(2x+1)}{12} + \frac{4(x-2)}{12}$		3	M1 for expressing both fractions correctly with a common denominator. Allow as two separate fractions.
		E.g. $\frac{6x+3+4x-8}{12}$			M1 for removing brackets correctly in a correct single fraction
			$\frac{10x-5}{12}$		A1 accept $\frac{5(2x-1)}{12}$

c	$(4^{k+3} =)(2^2)^{k+3}$ oe or $(16 =)2^4$ $(16 =)4^2$ or $(2^k =)\left(4^{\frac{1}{2}}\right)^k$ oe $(4^{k+3} =)\left(16^{\frac{1}{4}}\right)^{k+3}$ oe or $(2^k =)\left(16^{\frac{1}{4}}\right)^k$ oe		4	M1 for $(2^2)^{k+3}$ oe or 2^4 or 4^2 or $\left(4^{\frac{1}{2}}\right)^k$ oe or $\left(16^{\frac{1}{4}}\right)^{k+3}$ oe or $\left(16^{\frac{1}{4}}\right)^k$ oe
	$(4^{k+3} =)(2^2)^{k+3}$ oe and $(16 =)2^4$ $(16 =)4^2$ and $(2^k =)\left(4^{\frac{1}{2}}\right)^k$ oe $(4^{k+3} =)\left(16^{\frac{1}{4}}\right)^{k+3}$ oe and $(2^k =)\left(16^{\frac{1}{4}}\right)^k$ oe			M1 for $(2^2)^{k+3}$ oe and 2^4 or 4^2 and $\left(4^{\frac{1}{2}}\right)^k$ oe or $\left(16^{\frac{1}{4}}\right)^{k+3}$ oe and $\left(16^{\frac{1}{4}}\right)^k$ oe
	E.g. $2k + 6 = 4 + k$ or $k + 3 = 2 + \frac{1}{2}k$ or $\frac{1}{2}(k + 3) = 1 + \frac{1}{4}k$			M1 for a correct linear equation in k
		-2		A1 dep on at least M2
				Total 9 marks

13	e.g. $\begin{pmatrix} 5 \\ 3 \end{pmatrix} - \begin{pmatrix} -2 \\ 4 \end{pmatrix}$ or $\begin{pmatrix} 5 \\ 3 \end{pmatrix} + \begin{pmatrix} 2 \\ -4 \end{pmatrix}$		2	M1 or for $\begin{pmatrix} 7 \\ a \end{pmatrix}$ where $a \neq -1$ or $\begin{pmatrix} b \\ -1 \end{pmatrix}$ where $b \neq 7$
		$\begin{pmatrix} 7 \\ -1 \end{pmatrix}$		A1
				Total 2 marks

14	$BFD = 39^\circ$	$BED = 39^\circ$		4	B1
	$BDE = 180 - (18 + 39)$	$EBD = 18^\circ$ and $BDE = 180 - (18 + 39)$			M1
			123		A1
					B1 dep on M1 for all correct circle theorems relevant for their method e.g. <u>alternate segment theorem and opposite angles</u> in a <u>cyclic quadrilateral</u> sum to 180° or <u>alternate segment theorem and angles in same segment</u> are equal
					Total 4 marks

<p>15 a</p>	<p>E.g. $x = 4.57\dots$ and $100x = 457.57\dots$ or $10x = 45.757\dots$ and $1000x = 4575.7\dots$ or $x = 0.57\dots$ and $100x = 57.57\dots$ or $10x = 5.757\dots$ and $1000x = 575.7\dots$</p>		2	<p>M1 for selecting 2 recurring decimals that when subtracted give a whole number or terminating decimal eg 453 or 4530 etc eg $100x = 457.57\dots$ and $x = 4.57\dots$ or $1000x = 4575.7\dots$ and $10x = 45.757\dots$ with intention to subtract. (If recurring dots not shown then allow $10x = 45.757$, $100x = 457.57$, and $1000x = 4575.7$ to at least 5sf) or $4 + 0.5757$ and eg $x = 0.57\dots$, $100x = 57.57\dots$ with intention to subtract.</p>
	<p>E.g. $100x - x = 457.57\dots - 4.57\dots = 453$ and $\frac{453}{99} = \frac{151}{33}$ or $4\frac{19}{33}$ or $1000x - 10x = 4575.7\dots - 45.757\dots = 4530$ and $\frac{4530}{990} = \frac{151}{33}$ or $4\frac{19}{33}$ or $100x - x = 57.57\dots - 0.57\dots = 57$ and $\frac{57}{99}$ or $\frac{19}{33}$ (so) $4.\dot{5}\dot{7} = 4\frac{19}{33}$ or $1000x - 10x = 575.7\dots - 5.757\dots = 570$ and $\frac{570}{990}$ or $\frac{57}{99}$ or $\frac{19}{33}$ (so) $4.\dot{5}\dot{7} = 4\frac{19}{33}$</p>	Shown		<p>A1 for completion to $\frac{151}{33}$ or $4\frac{19}{33}$</p>

15 b	E.g. $\frac{2}{6-3\sqrt{2}} \times \frac{6+3\sqrt{2}}{6+3\sqrt{2}} \text{ or}$ $\frac{2}{6-3\sqrt{2}} \times \frac{-6-3\sqrt{2}}{-6-3\sqrt{2}}$		3	M1 for rationalising the denominator by multiplying numerator and denominator by $6+3\sqrt{2}$ (or $-6-3\sqrt{2}$)
	$\frac{12+6\sqrt{2}}{36-18\sqrt{2}+18\sqrt{2}-18} \text{ or}$ $\frac{12+6\sqrt{2}}{18} \text{ or } \frac{12+6\sqrt{2}}{6^2-(3\sqrt{2})^2} \text{ or } \frac{12+6\sqrt{2}}{6^2-9 \times 2}$			M1 (numerator may be expanded or denominator may be 4 terms which need to be all correct)
		$\frac{2+\sqrt{2}}{3}$		A1 or for stating $a = 2$ and $b = 3$ for $\frac{2+\sqrt{2}}{3}$ from correct working dep on M2
				Total 5 marks

16 a	E.g. $x^2 + 4x - 2x - 8 (= x^2 + 2x - 8)$ or $x^2 - 2x + x - 2 (= x^2 - x - 2)$ or $x^2 + 4x + x + 4 (= x^2 + 5x + 4)$		3	M1 for multiplying out two brackets correctly with no more than one error
	E.g. $x^3 + 2x^2 - 8x + x^2 + 2x - 8$ or $x^3 + 4x^2 - 2x^2 - 8x + x^2 + 4x - 2x - 8$ or $x^3 - x^2 - 2x + 4x^2 - 4x - 8$ or $x^3 - 2x^2 + x^2 - 2x + 4x^2 - 8x + 4x - 8$ or $x^3 + 5x^2 + 4x - 2x^2 - 10x - 8$ or $x^3 + 4x^2 + x^2 + 4x - 2x^2 - 8x - 2x - 8$			M1 for at least 3 terms correct out of a maximum of 6 terms or for at least 4 terms correct out of a maximum of 8 terms
		$x^3 + 3x^2 - 6x - 8$		A1
b	E.g. $(x - 5)^2 - 5^2 (+ 40)$ or $(x - 5)^2 - 25 (+ 40)$ $(x^2 + 2ax + a^2 (+ b^2))$ $2a = -10$ or $a = -5$		2	M1 for a correct first step or for equating coefficients
		$(x - 5)^2 + 15$		A1 accept $a = -5, b = 15$ SC B1 for $(-x + 5)^2 + 15$ or $(5 - x)^2 + 15$
				Total 5 marks

17	$y(6y + 5) - 2y^2 = 6$	$x\left(\frac{x-5}{6}\right) - 2\left(\frac{x-5}{6}\right)^2 = 6$		5	M1 for substitution of linear equation into quadratic or multiplying linear equation by y e.g. $xy - 6y^2 = 5y$ and intention to subtract the two equations
	E.g. $4y^2 + 5y - 6 (= 0)$ oe $4y^2 + 5y = 6$	E.g. $4x^2 - 10x - 266 (= 0)$ oe $4x^2 - 10x = 266$			A1 (dep on M1) writing the correct quadratic expression in form $ax^2 + bx + c (= 0)$ allow $ax^2 + bx = c$
	E.g. $(4y - 3)(y + 2) (= 0)$ $(y =) \frac{-5 \pm \sqrt{5^2 - 4 \times 4 \times -6}}{2 \times 4}$ $4 \left[\left(y + \frac{5}{8} \right)^2 - \left(\frac{5}{8} \right)^2 \right] = 6$ oe	E.g. $(2x - 19)(x + 7) (= 0)$ $(x =) \frac{5 \pm \sqrt{(-5)^2 - 4 \times 2 \times (-133)}}{2 \times 2}$ $4 \left[\left(x - \frac{10}{8} \right)^2 - \left(\frac{10}{8} \right)^2 \right] = 266$ oe			M1 (dep on M1) for a complete method to solve their 3-term quadratic equation (allow one sign error and some simplification – allow as far as $\frac{-5 \pm \sqrt{25 + 96}}{8}$ or $\frac{5 \pm \sqrt{25 + 1064}}{4}$)
	$(y =) \frac{3}{4}$ and $(y =) -2$	$(x =) \frac{19}{2}$ and $(x =) -7$			A1 Dep on first M1 for having two correct x values or two correct y values
			$x = \frac{19}{2}, y = \frac{3}{4}$ $x = -7, y = -2$		A1 Dep on first M1 Must be paired and labelled correctly
					Total 5 marks

18	E.g. $28 \div 2 (= 14)$ or $1\text{cm}^2 = 2$ students		5	M1 for method to find the frequency density for the first bar or any correct value on the fd axis or can be implied by a correct frequency (30 or 24 or 36)
	$2 \times 20 (= 40)$ $1 \times 30 (= 30)$ $1 \times 24 (= 24)$ $3 \times 12 (= 36)$ or 40, 30, 24, 36			M1 for method to find the missing frequencies (at least 3 correct)
	$1 \times 28 + 3 \times '40' + 4.5 \times '30' + 5.5 \times '24' + 7.5 \times '36' (= 685)$ or $28 + 120 + 135 + 132 + 270 (= 685)$			M1 (indep ft) for a method to find the total (mid value \times frequency) for at least 4 products using their values in the table (need not be evaluated) Allow consistent use of end points for at least 4 products which must be added
	' 685 ' \div $(28 + '40' + '30' + '24' + '36')$ $(= 4.335\dots)$ or ' 685 ' \div 158 $(= 4.335\dots)$			M1 (dep on previous M1)
		4.34		A1 accept 4.33 - 4.34
				Total 5 marks

19	7.75, 7.85, 3.35, 3.45, 13.5, 14.5		3	<p>B1 for sight of a correct upper or lower bound</p> <p>Accept</p> <p>$3.44\dot{9}$ for 3.45 or</p> <p>$7.84\dot{9}$ for 7.85 or</p> <p>$14.4\dot{9}$ for 14.5</p>
	$(k =) \frac{13.5}{7.85 - 3.35}$			<p>M1 for correct substitution into</p> $k = \frac{t_{LB}}{a_{UB} - h_{LB}}$ <p>where $13.5 \leq t_{LB} < 14$ and</p> <p>$7.8 < a_{UB} \leq 7.85$ and</p> <p>$3.35 \leq h_{LB} < 3.4$</p>
		3		A1 accept 3.0
				Total 3 marks

20	$(v =) 3t^2 - 9 \times 2t + 33$		5	M1 for differentiating at least 2 terms correctly
	$(a =) 3 \times 2t - '18'$ or $(t =) -\frac{-18}{2 \times 3} \left(= \frac{18}{6} \right)$	$(v =) 3 \left[(t-3)^2 - (3)^2 \right] (+33) \text{ or}$ $(v =) 3 \left[(t-3)^2 - (3)^2 (+11) \right]$		M1 dep ft must be a two term linear equation or for the use of $(t =) -\frac{b}{2a}$ or for a correct first step for completing the square on at least a two term quadratic
	$6t - 18 = 0 \text{ or } t = 3$	$(v =) 3 \left[(t-3)^2 - (3)^2 \right] + 33 \text{ or}$ $(v =) 3 \left[(t-3)^2 - (3)^2 + 11 \right]$		M1 dep on at least M2 for equating their acceleration to 0 or for a correct method for completing the square on at least a two term quadratic
	$3 \times '3'^2 - 18 \times '3' + 33$	$(v =) 3(t-3)^2 + 6 \text{ or}$ $(v =) 3 \left[(t-3)^2 + 2 \right]$		M1 dep on at least M2 for substituting their t into v or for a seeing a correct simplified expression after completing the square
			6	A1
				Total 5 marks

21	E.g. $a + 3d = 6$ oe		6	M1 for forming an equation for the 4 th term of the sequence
	E.g. $\frac{11}{2}(2a + 10d) = (a + 5d)^2 + 18$			M1 for forming an equation for the sum of the first 11 terms of the sequence
	E.g. $a = 6 - 3d$ and $\frac{11}{2}[2(6 - 3d) + 10d] = (6 - 3d + 5d)^2 + 18$ or $d = \frac{6 - a}{3}$ and $\frac{11}{2}\left[2a + 10\left(\frac{6 - a}{3}\right)\right] = \left(a + 5\left(\frac{6 - a}{3}\right)\right)^2 + 18$			M1 dep on M2 for a correct first step to solve the two equations (writing the equation in terms of one variable) Note: If $\frac{11}{2}(2a + 10d) = (a + 5d)^2 + 18$ is expanded then this must be a correct expansion E.g. $11a + 55d = a^2 + 10ad + 25d^2 + 18$
	E.g. $2d^2 + d - 6(=0)$ oe or $2a^2 - 27a + 36(=0)$ oe			A1 for a correct 3 term quadratic equation
	$d = 1.5$ oe and $a = 1.5$ oe			A1 for a correct value of d and a
		30		A1 cao
				Total 6 marks

22	$\left(\frac{-1+2}{2}, \frac{5+10}{2}\right)$ or (0.5, 7.5) oe		5	M1
	$\frac{10-5}{2-(-1)}\left(=\frac{5}{3}\right)$ oe			M1
	$m \times \frac{5}{3} = -1$ oe or $m = -\frac{3}{5}$ oe			M1 ft their gradient for use of $m_1 \times m_2 = -1$
	'7.5' = $-\frac{3}{5}$ ' × '0.5' + c or c = 7.8 oe or $y - '7.5' = -\frac{3}{5}(x - '0.5')$			M1 ft dep on first M1 and third M1
		$5y + 3x = 39$		A1 oe where p , q and r must be integers
				Total 5 marks