

Mark Scheme (Results)

January 2016

Pearson Edexcel International GCSE
Mathematics A (4MA0)
Paper 3H

Pearson Edexcel Certificate
Mathematics A (KMA0)
Paper 3H

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2016

Publications Code UG043259

All the material in this publication is copyright

© Pearson Education Ltd 2016

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- **Types of mark**
 - M marks: method marks
 - A marks: accuracy marks
 - B marks: unconditional accuracy marks (independent of M marks)
- **Abbreviations**
 - cao – correct answer only
 - ft – follow through
 - isw – ignore subsequent working
 - SC - special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - eoo – each error or omission
 - awrt –answer which rounds to

- **No working**

If no working is shown then correct answers normally score full marks

If no working is shown then incorrect (even though nearly correct) answers score no marks.
- **With working**

If there is a wrong answer indicated on the answer line always check the working in the body of the script (and on any diagrams), and award any marks appropriate from the mark scheme.

If it is clear from the working that the "correct" answer has been obtained from incorrect working, award 0 marks.

Any case of suspected misread loses A (and B) marks on that part, but can gain the M marks.

If working is crossed out and still legible, then it should be given any appropriate marks, as long as it has not been replaced by alternative work.

If there is a choice of methods shown, then no marks should be awarded, unless the answer on the answer line makes clear the method that has been used.

If there is no answer on the answer line then check the working for an obvious answer.
- **Ignoring subsequent work**

It is appropriate to ignore subsequent work when the additional work does not change the answer in a way that is inappropriate for the question: eg. Incorrect cancelling of a fraction that would otherwise be correct.

It is not appropriate to ignore subsequent work when the additional work essentially makes the answer incorrect eg algebra.

Transcription errors occur when candidates present a correct answer in working, and write it incorrectly on the answer line; mark the correct answer.
- **Parts of questions**

Unless allowed by the mark scheme, the marks allocated to one part of the question CANNOT be awarded in another.

International GCSE Maths Jan 2016 – Paper 3H Mark scheme				
Apart from Questions 8, 15c, 16 and 22 (where the mark scheme states otherwise), the correct answer, unless clearly obtained by an incorrect method, should be taken to imply a correct method.				
Q	Working	Answer	Mark	Notes
1 a	eg. $24 \div 6 \times 800$			M1 for a complete method
		3200	2	A1 accept 3.2 litres
b	eg. $450 \div 300 \times 6$			M1 for a complete method
		9	2	A1
				Total 4 marks

2	1.75 or $1\frac{3}{4}$ or 105			M1 for correctly converting "1hr 45 mins" into a decimal or fraction or minutes (eg. 1.75h or 105 min)
	$140 \div$ "1.75" "or $\frac{140}{105} \times 60$			M1 independent but "1.75" or "105" must be correct for their duration accept $140 \div 1.45$
		80	3	A1
				Total 3 marks

3	$\frac{3}{8} \times \frac{12}{7}$			M1
		$\frac{36}{56}$ oe	2	A1 dep on M1 Accept $\frac{9}{14}$ if clear cancelling seen NB: Use of decimals gains M0 A0
	Alternative : $\frac{9n}{24n} \div \frac{14n}{24n}$ for any integer n			M1 Must see an intention to divide
		$\frac{9}{14}$ oe	2	A1 dep on M1 Answer must come directly from their method eg. $\frac{36}{96} \div \frac{56}{96}$ must be followed by $\frac{36}{56}$
				Total 2 marks

4	a		$5(3r + 2)$	1	B1
	b		y^9	1	B1
	c	$x^2 + 5x - x - 5$			M1 for 3 correct terms out of a maximum of 4 terms or for 4 correct terms ignoring signs or for $x^2 + 4x + k$ for any non-zero value of k or for $\dots + 4x - 5$
			$x^2 + 4x - 5$	2	A1
	d				M1 for any one correct application of index laws in a product or quotient eg. $\frac{36m^4}{30k^2m}$, $\frac{36k^3m^3}{30k^5}$
			$\frac{6m^3}{5k^2}$	2	A1 oe eg. $1.2m^3k^{-2}$
					Total 6 marks

5	a		20 to 24 oe	1	B1
	b	$2 \times 6 + 7 \times 3 + 12 \times 5 + 17 \times 12 + 22 \times 14$ or $12 + 21 + 60 + 204 + 308$ or 605			M2 freq \times all correct midpoint values stated (or evaluated) with intention to add (condone any two errors in midpoints or frequencies) If not M2 then award M1 for all products $t \times f$ (and t is consistently within the interval, including end values) and intention to add (condone any two errors in their midpoints or frequencies)
		"605" \div 40 or $\frac{2 \times 6 + 7 \times 3 + 12 \times 5 + 17 \times 12 + 22 \times 14}{6 + 3 + 5 + 12 + 14}$			M1 dep on at least M1
			15.125	4	A1 accept 15 or 15.1 or 15.13 from correct working with no errors (15 without working gains M0 A0) NB. Accept 15.1625 (using 2.25 as mi-interval in first class)
	c	$14 \div 40 \times 100$ oe			M1 award M1 for $26 \div 40 \times 100$ or 65%
			35	2	A1
					Total 7 marks

6		bisector with construction arcs	2	B2 for bisector within guidelines with two pairs of relevant construction arcs seen If not B2 then B1 for a bisector within guidelines with no arcs present or relevant arcs present with no bisector
				Total 2 marks

7	a	1,2,3,4,5,6,7,9	1	B1 no repeats
	b	eg. 4, 5, 7, 8	2	B2 for 4, 5, and any two of 7, 8, 9, 10 If not B2 then B1 for 4,5 or 4,5 and any one or two other numbers from the universal set or 4, 5, 7, 8, 9, 10 or 7, 8, 9, 10 or Venn diagram with 4,5 in intersection and any two correct numbers (eg. 8,10) in C
				Total 3 marks

8	$6x + 15$			M1 for correct expansion of bracket OR division of all terms in a correct equation by 3
	$6x + x = 4 - 15$			M1 for correct rearrangement within a correct equation with x terms on one side and numbers on the other
		$-1\frac{4}{7}$ oe	3	A1 Award full marks for a correct answer if at least 1 method mark awarded (allow $-\frac{11}{7}$ as final answer) accept $-1.57(1428\dots)$
				Total 3 marks

9	a	$\pi \times 5.4^2 \times 16$			M1
			1466	2	A1 answer in range 1464.9 - 1466
	bi		5.45	1	B1 accept 5.449
	bii		5.35	1	B1
					Total 4 marks

10	a	(-2, -1) (-1, 1) (0, 3) (1, 5) (2, 7) (3, 9) (4, 11)	correct line	3	B3 for $y = 2x + 3$ drawn from $x = -2$ to 4 if not B3 then B2 for a correct straight line segment through at least 3 of (-2, -1) (-1, 1) (0, 3) (1, 5) (2, 7) (3, 9) (4, 11) OR for all of (-2, -1) (-1, 1) (0, 3) (1, 5) (2, 7) (3, 9) (4, 11) plotted but not joined if not B2 then B1 for any 2 correct points stated (could be in a table) or plotted OR a line with a positive gradient through (0, 3) OR a line with gradient 2
	b				M1 for $x = 3$ and $y = 2$ drawn
			correct region	2	A1 for correct region identified (R need not be labelled) Accept shaded or unshaded
					Total 5 marks

11	$\tan ACB = \frac{4.5}{9.6}$			M1 for correct trig statement eg. $\sin ACB = \frac{4.5}{\sqrt{112.41}}$ or $\cos ACB = \frac{9.6}{\sqrt{112.41}}$	
	$\tan^{-1} \left(\frac{4.5}{9.6} \right)$			M1 dep	
		25.1	3	A1 awrt 25.1	
					Total 3 marks

12	$5t - 5g = 2t + 7$			M1 for expanding bracket within the equation or division of all terms by 5
	$5t - 2t = 7 + 5g$			M1 (ft a 4 term equation) to isolate terms in t
		$t = \frac{5g + 7}{3}$	3	A1 oe
				Total 3 marks

13	a	11 12 <u>13</u> 13 16 17 17 18 <u>19</u> 19 20			M1 13 or 12.75 (LQ) 19 or 18.25 (UQ) identified from ordered list OR attempt to find IQR eg. 3(rd) and 9(th) seen or 2.75(th) and 8.25(th) seen
		11 12 <u>13</u> 13 16 <u>17</u> 17 18 <u>19</u> 19 20 or 19 - 13			M1 Identify 13 or 12.75 (LQ), AND 19 or 18.25 (UQ)
			6	3	A1 accept 5.5
	b		James and reason using IQR	1	B1 ft from (a) James - he has a lower IQR oe (IQR must be part of the statement)
	c		no change with reason	1	B1 no change box ticked with reason eg. 2 new scores above median and 2 new scores below median or median of 4 numbers is 17
Total 5 marks					

14 a	8000×1.045 oe (=8360)			M1 or 8000×1.0275^3 (=8678.316375)	M2 for $8000 \times 1.045 \times 1.0275^3$
	"8360" $\times 1.0275^3$ oe			M1 "8678.316375" $\times 1.045$	
			9068.84	3	A1 accept 9069 and answers in the range 9068.8(0) - 9068.9(0) SC: B1 for an answer of 9020 ($8000 + 360 + 3 \times 220$)
b	$1 + 0.02$ (=1.02)			M1	M1 for $100(\%) + 2(\%)$ (= 102(%))
	$5763 \div "1.02"$ oe			M1 dep	M1 (dep) for $5763 \div "102" \times 100$ oe
			5650	3	A1
					Total 6 marks

15 a		5	1	B1
b		$(-7 \leq) k < 5,$ $13 < k (\leq 25)$	2	B2 accept $k \leq 5, k \geq 13$ If not B2 then B1 for 5 and 13 or $k < 5$ or $k > 13$
c	tangent drawn at $x = -2.5$			M1
		7.5	3	M1 complete method to find gradient A1 accept answers in range 7 - 8 with working seen
d		-2	1	B1
e	$\frac{1}{2+3}$ or -1 or $f(-1)$			M1
		9	2	A1
Total 9 marks				

16	$x^2 < 9$ or $x^2 - 9 < 0$ or $\frac{\pm\sqrt{-4 \times 5 \times -45}}{2 \times 5}$			M1 Allow $x^2 = 9$ or $x^2 - 9 = 0$
	$\pm\sqrt{9}$ or ± 3 or $(x + 3)(x - 3)$ or $x < 3$ or $x > -3$			M1
		$-3 < x < 3$	3	A1
				Total 3 marks

17	ai		96	1	B1
	aii		<u>Angle at the centre is twice angle at the circumference</u>	1	B1 (indep)
b		$73 - 26$			M1 for a complete method
			47		A1
				3	B1 (dep on M1) <u>Alternate segment theorem</u>
b	Alternative Scheme				
		Angle $RST = 180 - 73 (=107)$ and Angle $SRT = 180 - 26 - 107$			M1
			47		A1
					B1 (dep on M1) <u>Alternate segment theorem</u>
					Total 5 marks

19	$\frac{1}{3}\pi(3r)^2h$			M1 or $\frac{4}{3}\pi\left(\frac{y}{3}\right)^3$ Condone missing brackets around $3r$ or $\frac{y}{3}$ for this mark only Allow use of 3.14 consistently (or other correct value)
	$\frac{4}{3}\pi r^3 = \frac{1}{3}\pi(3r)^2h$			M1 or $\frac{4}{3}\pi\left(\frac{y}{3}\right)^3 = \frac{1}{3}\pi y^2h$
		$\frac{9h}{4}$ oe	3	A1 for $\frac{9h}{4}$ oe or $r = \frac{9h}{4}$ NB : $h = \frac{4r}{9}$ scores M2 A0
				Total 3 marks

20	$\frac{\sin A}{36} = \frac{\sin 48}{57}$			M1 or $\frac{36}{\sin A} = \frac{57}{\sin 48}$
	$A = \sin^{-1}\left(\frac{\sin 48}{57} \times 36\right)$ or A in range 27.9 – 28			M1 dep
	$\frac{1}{2} \times 57 \times 36 \times \sin(180 - 48 - "28")$ (=995.49...)			M1 dep on the first M1 or $\frac{1}{2} \times AC \times 36 \times \sin(48)$ with AC in range 74 – 74.5 or AC from a correct method
		995	4	A1 for answer in the range 995 - 996
	Alternative			
	Let BD be altitude of triangle $BD = 36 \times \sin 48^\circ$ (= 26.7..) AND $DC = 36 \times \cos 48^\circ$ (= 24.0..)			M1 correct method to find BD and DC
	$AD = \sqrt{57^2 - "26.7.."^2} = 50.331$			M1 dep
	$\frac{1}{2} \times "26.7.." \times ("24.0.." + "50.331")$			M1 dep on the first M1
		995	4	A1 for answer in the range 995 - 996
				Total 4 marks

21	Scheme 1 $P(L,L) + P(L, NL) + P(NL, L)$			Bus L = 0.07 ; BusNL = 0.63 BikeL = 0.015; BikeNL = 0.285 correct products may be seen instead of these values
	$1 - 0.1 (=0.9)$ or $1 - 0.05 (=0.95)$ or $1 - 0.7 (=0.3)$			M1
	P(not late for work) $= 0.7 \times 0.9 + 0.3 \times 0.95 (= 0.915 \text{ or } \frac{183}{200})$			M1 or at least 4 correct products from (addition not needed) $0.07^2 + 2 \times 0.07 \times 0.63 + 2 \times 0.07 \times 0.015 +$ $2 \times 0.07 \times 0.285 + 2 \times 0.63 \times 0.015 + 0.015^2 +$ $2 \times 0.015 \times 0.285$
	P(late for work) $= 0.7 \times 0.1 + 0.3 \times 0.05 (= 0.085 \text{ or } \frac{17}{200})$ OR $1 - "0.915" (=0.085)$			M1 or at least 8 correct products from (addition not needed) $0.07^2 + 2 \times 0.07 \times 0.63 + 2 \times 0.07 \times 0.015 +$ $2 \times 0.07 \times 0.285 + 2 \times 0.63 \times 0.015 + 0.015^2 +$ $2 \times 0.015 \times 0.285$
	P(late on at least one day) $= 2 \times "0.085" \times "0.915" + "0.085" \times "0.085"$			M1 or all 12 correct products with addition : dep $0.07^2 + 2 \times 0.07 \times 0.63 + 2 \times 0.07 \times 0.015 +$ $2 \times 0.07 \times 0.285 + 2 \times 0.63 \times 0.015 + 0.015^2 +$ $2 \times 0.015 \times 0.285$
	0.162775 or $\frac{6511}{40000}$ oe	5	A1	
				Total 5 marks

21	Scheme 2 $1 - P(\text{NL}, \text{NL})$			BusNL = 0.63 BikeNL = 0.285 correct products may be seen instead of these values
	$1 - 0.1 (=0.9)$ or $1 - 0.05 (=0.95)$ or $1 - 0.7 (=0.3)$			M1
	P(not late for work) $= 0.7 \times 0.9 + 0.3 \times 0.95 (= 0.915 \text{ or } \frac{183}{200})$			M1 or any two correct products from (addition not needed) $0.63^2 + 0.285^2 + 2 \times 0.63 \times 0.285$
	P(not late both days) $= "0.915" \times "0.915" (= 0.837225)$			M1 or all correct products dep (addition not needed) $0.63^2 + 0.285^2 + 2 \times 0.63 \times 0.285$ $(= 0.3969 + 0.081225 + 2 \times 0.17955 = 0.837225)$
	P(late on at least one day) $= 1 - "0.837225" = 0.162775$			M1 dep
	0.162775 or $\frac{6511}{40000}$ oe	5	A1	
				Total 5 marks

22	(length =) $\frac{28-2x}{2}$ or $14-x$ or $\sqrt{12^2-x^2}$			M1 correct expression for length of rectangle OR a pair of correct simultaneous equations eg. $x^2 + y^2 = 12^2$ and $2x + 2y = 28$
	$12^2 = x^2 + (14-x)^2$ oe			M1 for correct equation in one variable accept other forms eg. $2\sqrt{12^2-x^2} + 2x = 28$
	$144 = x^2 + 196 - 28x + x^2$ or $144 = x^2 + \frac{784-112x+4x^2}{4}$			M1 (indep) for expansion of brackets $196 - 28x + x^2$ or $\frac{784-112x+4x^2}{4}$
	$2x^2 - 28x + 52 = 0$ or $x^2 - 14x + 26 = 0$			A1 for a correct simplified quadratic equation
	eg. $x = \frac{- -14 \pm \sqrt{(-14)^2 - 4 \times 1 \times 26}}{2 \times 1}$			M1 ft for correct substitution into quadratic formula for their quadratic (condone one sign error; condone missing brackets and 14^2)
	eg. $\frac{14 \pm \sqrt{92}}{2}$ or $7 \pm \sqrt{23}$			M1 (indep) correct simplification of discriminant for correct quadratic equation
		11.8, 2.20	7	A1 answers in the ranges 11.7 - 11.8 and 2.2 - 2.21 dep on a correct quadratic equation and at least M4
				Total 7 marks

23	$\overrightarrow{OP} = 4\mathbf{a}$ or $\overrightarrow{PO} = 4\mathbf{a}$ or $\overrightarrow{PA} = 2\mathbf{a}$ or $\overrightarrow{AP} = -2\mathbf{a}$			M1 for correct use of ratio to find a relevant vector NB: may be seen on diagram
	$\overrightarrow{AB} = 4\mathbf{b} - 6\mathbf{a}$ oe or $\overrightarrow{AM} = 2\mathbf{b} - 3\mathbf{a}$ oe or $\overrightarrow{MB} = 2\mathbf{b} - 3\mathbf{a}$ oe			M1
	$\overrightarrow{PM} = 2\mathbf{b} - \mathbf{a}$ oe or $\overrightarrow{MC} = 6\mathbf{b} - 3\mathbf{a}$ oe or $\overrightarrow{PC} = 8\mathbf{b} - 4\mathbf{a}$ oe			M1 indep correct vector equation for \overrightarrow{PM} or \overrightarrow{MC} or \overrightarrow{PC} in terms of \mathbf{a} and/or \mathbf{b}
	$\overrightarrow{PM} = 2\mathbf{b} - \mathbf{a}$ oe AND $\overrightarrow{MC} = 6\mathbf{b} - 3\mathbf{a}$ oe or $\overrightarrow{PM} = 2\mathbf{b} - \mathbf{a}$ oe AND $\overrightarrow{PC} = 8\mathbf{b} - 4\mathbf{a}$ oe or $\overrightarrow{PC} = 8\mathbf{b} - 4\mathbf{a}$ oe AND $\overrightarrow{MC} = 6\mathbf{b} - 3\mathbf{a}$ oe			A1
		Show	5	A1 Completion with final reasons given using simplified vectors
				Total 5 marks

