Candidate Name _____ Centre Number | | | Candidate

International General Certificate of Secondary Education
CAMBRIDGE INTERNATIONAL EXAMINATIONS

CHEMISTRY 0620/3

PAPER 3

OCTOBER/NOVEMBER SESSION 2002

1 hour 15 minutes

Candidates answer on the question paper.
No additional materials are required.

TIME 1 hour 15 minutes

INSTRUCTIONS TO CANDIDATES

Write your name, Centre number and candidate number in the spaces at the top of this page.
Answer **all** questions.
Write your answers in the spaces provided on the question paper.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets [ ] at the end of each question or part question.
A copy of the Periodic Table is printed on page 12.

FOR EXAMINER'S USE	
1	
2	
3	
4	
5	
TOTAL	

This question paper consists of 10 printed pages and 2 blank pages.

UNIVERSITY of CAMBRIDGE
Local Examinations Syndicate
[Turn over

1 **(a)** Sulphuric acid is made by the Contact Process.

$$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g) \quad \text{forward reaction is exothermic}$$

(i) What are the reaction conditions for the Contact Process?

.....

.....[3]

(ii) Would the yield of sulphur trioxide increase, decrease or stay the same when the temperature is increased? Explain your answer.

.....

.....

.....[2]

(iii) Describe how sulphur trioxide is changed into concentrated sulphuric acid.

.....

.....[2]

(b) There are three ways of making salts from sulphuric acid.
titration using a burette and indicator
precipitation by mixing the solutions and filtering
neutralisation of sulphuric acid using an excess of an insoluble base

Complete the following table of salt preparations.

method	reactant 1	reactant 2	salt
titration	sulphuric acid		sodium sulphate
neutralisation	sulphuric acid		zinc sulphate
precipitation	sulphuric acid		barium sulphate
	sulphuric acid	copper(II) oxide	copper(II) sulphate

[4]

(c) The results of an investigation into the action of heat on copper(II) sulphate-5-water, a blue crystalline solid, are given below.

The formula is $CuSO_4.5H_2O$ and the mass of one mole is 250 g

A 5.0 g sample of the blue crystals is heated to form 3.2 g of a white powder. With further heating this decomposes into a black powder and sulphur trioxide.

(i) Name the white powder.

.....[1]

(ii) What is observed when water is added to the white powder?

.....[1]

(iii) Name the black powder.

.....[1]

(iv) Calculate the mass of the black powder. Show your working.

.....

.....

.....[3]

2 Manganese is a transition element. It has more than one valency and the metal and its compounds are catalysts.

(a) **(i)** Predict **three** other properties of manganese that are typical of transition elements.

.....

.....[3]

(ii) Complete the electron distribution of manganese by inserting one number.

2 + 8 + ... + 2 [1]

(b) It has several oxides, three of which are shown below.
Manganese(II) oxide, which is basic.
Manganese(III) oxide, which is amphoteric.
Manganese(IV) oxide, which is acidic.

(i) Complete the word equation.

manganese(II) oxide + hydrochloric acid → ..... + ... + ...

[2]

(ii) Which, if any, of these oxides will react with sodium hydroxide?

.....[1]

[Turn over

(c) Aqueous hydrogen peroxide decomposes to form water and oxygen.

$$2H_2O_2(aq) \rightarrow 2H_2O(l) + O_2(g)$$

This reaction is catalysed by manganese(IV) oxide

The following experiments were carried out to investigate the rate of this reaction.

A 0.1 g sample of manganese(IV) oxide was added to 20 cm^3 of 0.2 M hydrogen peroxide solution. The volume of oxygen produced was measured every minute. The results of this experiment are shown on the graph.



(i) How does the rate of reaction vary with time? Explain why the rate varies.

.....

.....[3]

(ii) The following experiment was carried out at the same temperature.

0.1 g of manganese(IV) oxide and 20 cm^3 of 0.4 M hydrogen peroxide

Sketch the curve for this experiment on the same grid. [2]

(iii) How would the shape of the graph differ if only half the mass of catalyst had been used in these experiments?

.....

.....

.....[2]

3 The elements in Period 3 and some of their common oxidation states are shown below.

Element	Na	Mg	A*l*	Si	P	S	C*l*	Ar
Oxidation State	+1	+2	+3	+4	–3	–2	–1	0

(a) **(i)** Why do the oxidation states increase from sodium to silicon?

.....[1]

(ii) After Group(IV) the oxidation states are negative and decrease across the period. Explain why.

.....

.....[2]

(b) The following compounds contain two elements. Predict their formulae.

aluminium sulphide .....

silicon phosphide ..... [2]

(c) Choose a different element from Period 3 that matches each description.

(i) It has a similar structure to diamond.

.....[1]

(ii) It reacts violently with cold water to form a solution pH = 14.

.....[1]

(iii) It has a gaseous oxide of the type XO_2, which is acidic.

.....[1]

(d) The only oxidation state of argon is zero. Why it is used to fill light bulbs?

.....

.....[1]

(e) Draw a diagram that shows the arrangement of the valency electrons in the ionic compound sodium phosphide.

Use o to represent an electron from sodium.
Use x to represent an electron from phosphorus. [3]

(f) Sodium reacts with sulphur to form sodium sulphide.

$$2Na + S \rightarrow Na_2S$$

An 11.5 g sample of sodium is reacted with 10 g of sulphur. All of the sodium reacted but there was an excess of sulphur.
Calculate the mass of sulphur left unreacted.

(i) Number of moles of sodium atoms reacted = ...
[2 moles of Na react with 1 mole of S]

(ii) Number of moles of sulphur atoms that reacted = ...

(iii) Mass of sulphur reacted = ...g

(iv) Mass of sulphur left unreacted = ...g [4]

4 For over 5000 years copper has been obtained by the reduction of its ores. More recently the metal has been purified by electrolysis.

(a) Copper is used to make alloys.

(i) Give **two** other uses of copper.

.....[2]

(ii) Alloys have similar structures to pure metals. Give a labelled diagram that shows the structure of a typical alloy, such as brass.

[3]

(b) Copper is refined by the electrolysis of aqueous copper(II) sulphate using copper electrodes. Describe the change that occurs at the electrodes.

(i) cathode (pure copper) .....

.....[1]

(ii) anode (impure copper) .....

.....[1]

(iii) Write an ionic equation for the reaction at the cathode.

.....[1]

(iv) If carbon electrodes are used, a colourless gas is given off at the anode and the electrolyte changes from a blue to a colourless solution.

The colourless gas is ..... .

The solution changes into ..... . [2]

(c) Electrolysis and cells both involve chemical reactions and electricity.

What is the essential difference between them?

.....

.....[2]

(d) Copper is an unreactive metal. Its compounds are easily reduced to the metal or decomposed to simpler compounds. Complete the following equations.

(i) ...CuO + ... $\rightarrow$ ...Cu + ...

(ii) Copper(II) hydroxide $\xrightarrow[\text{(heat)}]{}$ ..... + ...

.....

(iii) $Cu(NO_3)_2 \xrightarrow[\text{(heat)}]{}$ ..... + ... + ...

[4]

[Turn over

5 Alkenes are unsaturated hydrocarbons. They show structural isomerism. Alkenes take part in addition reactions and form polymers.

(a) Structural isomers have the same molecular formula but different structural formulae. Give an example of structural isomerism.

molecular formula .....

two structural formulae

[3]

(b) Ethene reacts with each of the following. Give the name and structural formula of each product.

(i) steam

name of product .....

structure of product

[2]

(ii) hydrogen

name of product .....

structure of product

[2]

(c) Alkenes polymerise by addition.

(i) Explain the term *polymerise*.

.....

.....[2]

(ii) What is the difference between addition polymerisation and condensation polymerisation?

.....

.....[2]

(iii) Poly(dichloroethene) is used extensively to package food. Draw its structure. The structural formula of dichloroethene is drawn below.



[2]

(d) Steel may be coated with another metal, eg zinc or chromium, or with a polymer, eg poly(chloroethene), to prevent rusting.

(i) Suggest a property of poly(chloroethene) that makes it suitable for this purpose.

.....[1]

(ii) Explain why the steel will rust when the protective coating of chromium or polymer is broken.

.....[1]

(iii) When the protective layer of zinc is broken, the steel still does not rust.
Suggest an explanation.

.....

.....

.....[2]

BLANK PAGE

BLANK PAGE

DATA SHEET
The Periodic Table of the Elements

Group																	
I	II										III	IV	V	VI	VII	0
							1 H Hydrogen 1									4 He Helium 2
7 Li Lithium 3	9 Be Beryllium 4										11 B Boron 5	12 C Carbon 6	14 N Nitrogen 7	16 O Oxygen 8	19 F Fluorine 9	20 Ne Neon 10
23 Na Sodium 11	24 Mg Magnesium 12										27 Al Aluminium 13	28 Si Silicon 14	31 P Phosphorus 15	32 S Sulphur 16	35.5 Cl Chlorine 17	40 Ar Argon 18
39 K Potassium 19	40 Ca Calcium 20	45 Sc Scandium 21	48 Ti Titanium 22	51 V Vanadium 23	52 Cr Chromium 24	55 Mn Manganese 25	56 Fe Iron 26	59 Co Cobalt 27	59 Ni Nickel 28	64 Cu Copper 29	65 Zn Zinc 30	70 Ga Gallium 31	73 Ge Germanium 32	75 As Arsenic 33	79 Se Selenium 34	80 Br Bromine 35	84 Kr Krypton 36
85 Rb Rubidium 37	88 Sr Strontium 38	89 Y Yttrium 39	91 Zr Zirconium 40	93 Nb Niobium 41	96 Mo Molybdenum 42	Tc Technetium 43	101 Ru Ruthenium 44	103 Rh Rhodium 45	106 Pd Palladium 46	108 Ag Silver 47	112 Cd Cadmium 48	115 In Indium 49	119 Sn Tin 50	122 Sb Antimony 51	128 Te Tellurium 52	127 I Iodine 53	131 Xe Xenon 54
133 Cs Caesium 55	137 Ba Barium 56	139 La Lanthanum 57 *	178 Hf Hafnium 72	181 Ta Tantalum 73	184 W Tungsten 74	186 Re Rhenium 75	190 Os Osmium 76	192 Ir Iridium 77	195 Pt Platinum 78	197 Au Gold 79	201 Hg Mercury 80	204 Tl Thallium 81	207 Pb Lead 82	209 Bi Bismuth 83	Po Polonium 84	At Astatine 85	Rn Radon 86
Fr Francium 87	226 Ra Radium 88	227 Ac Actinium 89 †															

140 Ce Cerium 58	141 Pr Praseodymium 59	144 Nd Neodymium 60	Pm Promethium 61	150 Sm Samarium 62	152 Eu Europium 63	157 Gd Gadolinium 64	159 Tb Terbium 65	162 Dy Dysprosium 66	165 Ho Holmium 67	167 Er Erbium 68	169 Tm Thulium 69	173 Yb Ytterbium 70	175 Lu Lutetium 71
232 Th Thorium 90	Pa Protactinium 91	238 U Uranium 92	Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	Bk Berkelium 97	Cf Californium 98	Es Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103

*58-71 Lanthanoid series
†90-103 Actinoid series

Key: a = relative atomic mass; X = atomic symbol; b = proton (atomic) number

The volume of one mole of any gas is 24 dm^3 at room temperature and pressure (r.t.p.).